



7B2 Monoclonal Antibody

Catalog No	YP-mAb-05831
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SCG5 SGNE1
Protein Name	Neuroendocrine protein 7B2 (Pituitary polypeptide) (Secretogranin V) (Secretogranin-5) (Secretory granule endocrine protein I) [Cleaved into: N-terminal peptide; C-terminal peptide]
Immunogen	Synthesized peptide derived from human protein . at AA range: 90-170
Specificity	7B2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	23kD
Cell Pathway	Secreted. Neuroendocrine and endocrine secretory granules.
Tissue Specificity	Brain,Pancreas,Pituitary,
Function	function:Acts as a molecular chaperone for PCSK2/PC2, preventing its premature activation in the regulated secretory pathway. Binds to inactive PCSK2 in the endoplasmic reticulum and facilitates its transport from there to later compartments of the secretory pathway where it is proteolytically matured and activated. Also required for cleavage of PCSK2 but does not appear to be involved in its folding. Plays a role in regulating pituitary hormone secretion. The C-terminal peptide inhibits PCSK2 in vitro.,PTM:Proteolytically cleaved in the Golgi by a furin-like convertase to generate bioactive peptides.,PTM:Sulfated on tyrosine residues.,similarity:Belongs to the 7B2 family.,subcellular location:Neuroendocrine and endocrine secretory granules.,subunit:Interacts with PCSK2/PC2 early in the secretory pathway. Dissociation occurs at later stages.,
Background	This gene encodes a secreted chaperone protein that prevents the aggregation of other secreted proteins, including proteins that are associated with neurodegenerative and metabolic disease. The encoded protein may be best



known for its role in the trafficking and activation of prohormone convertase PC2 (encoded by Gene ID: 5126). Phosphorylation of the encoded protein has been shown to have an inhibitory effect on its chaperone function. [provided by RefSeq, Jul 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using 7B2 Monoclonal Antibody