





PGK2 Monoclonal Antibody

Catalog No YP-mAb-05926 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name PGK2 PGKB Protein Name Phosphoglycerate kinase 2 (EC 2.7.2.3) (Phosphoglycerate kinase, testis specific) Immunogen Synthesized peptide derived from human protein . at AA range: 80-160 Specificity PGK2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (et protein level), or elderly men. Function Asialyic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glycerate in lavel phosphoglycerate kinase armity, astbunit (Monomer, Issues specificity). Testis spec		
Reactivity	Catalog No	YP-mAb-05926
Applications Gene Name PGK2 PGKB Protein Name Phosphoglycerate kinase 2 (EC 2.7.2.3) (Phosphoglycerate kinase, testis specific) Immunogen Synthesized peptide derived from human protein . at AA range: 80-160 Specificity PGK2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function Purity Purity Synonyms Cell Pathway Cytoplasm Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function Phosphate, online information:Phosphoglycerate in Apply phosphates, online information:Phosphoglycerate kinase entry, pathway. Carbohydrate degradation, glycoylsy, parturate from D-glycerate kinase farmily, subunit. Monomer, its sue specificity. Testis specific. Background Phosphoglycerate kinase 2(PGK2) Homo sapiens Tis gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate kinase in the later stages of	Isotype	IgG
Gene Name PGK2 PGKB Protein Name Phosphoglycerate kinase 2 (EC 2.7.2.3) (Phosphoglycerate kinase, testis specific) Immunogen Synthesized peptide derived from human protein . at AA range: 80-160 Specificity PGK2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroly phosphate. online information:Phosphoglycerate kinase entry pathway:Carbohydrate degradation: glycolysis: pyruvate from D-glyceraldehyde 3-phosphate: step 2/5. similarity Belongs to the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseu	Reactivity	Human;Mouse
Protein Name	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 80-160 Specificity PGK2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroly phosphate, online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5, similarity:Belongs to the phosphoglycerate kinase family , subunit:Monomer., tissue specificity Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosp	Gene Name	PGK2 PGKB
Specificity PGK2 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity.ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroly phosphate, online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5., similarity:Belongs to the phosphoglycerate kinase family, subunit Monomer., itssue specificity. Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the erooded protein is actu	Protein Name	Phosphoglycerate kinase 2 (EC 2.7.2.3) (Phosphoglycerate kinase, testis specific)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate, online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5. similarity:Belongs to the phosphoglycerate kinase family., subunit:Monomer., tissue specificity:Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embled-Meyerhof-Parmas pathway of glycolysis, in the later stages of	Immunogen	Synthesized peptide derived from human protein . at AA range: 80-160
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function Catallytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate. online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphoglycerate kinase family. subunit:Monomer.,tissue specificity:Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1.3-bisphosphoglycerate kinase that catalyzes the reversible conversion of 1.3-bisphosphoglycerate kinase in the later stages of the later stages of the phosphoglycerate kinase that catalyzes the reversible conversion of 1.3-bisphosphoglycerate kinase that catalyzes the reversible conversion of 1.3-bisphosphoglycerate kin	Specificity	PGK2 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphateonline information:Phosphoglycerate kinase entry.pathway:Carbohydrate degradation, glycoyls: pyruvate from D-glyceraldehyde 3-phosphate: step 2/5. similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate.	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm. Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate, online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5. similarity:Belongs to the phosphoglycerate kinase family., subunit:Monomer.,tissue specificity:Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis: Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm. Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate. online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate kinase so flosphosphoglycerate to 3-phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate.	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate kinase that catalyzes the reversible c	Dilution	WB 1:500-1:2000
Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific. Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Concentration	1 mg/ml
Synonyms Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry, pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Purity	≥90%
Observed Band 45kD Cell Pathway Cytoplasm . Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Synonyms	
Tissue Specificity Mainly found in round spermatids. Localized on the principle piece in the sperm (at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Observed Band	45kD
(at protein level). Testis-specific. Expression significantly decreased in the testis of elderly men. Function catalytic activity:ATP + 3-phospho-D-glycerate = ADP + 3-phospho-D-glyceroyl phosphate.,online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Cell Pathway	Cytoplasm .
phosphate.,online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis specific., Background phosphoglycerate kinase 2(PGK2) Homo sapiens This gene is intronless, arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Tissue Specificity	(at protein level). Testis-specific. Expression significantly decreased in the testis
arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of	Function	phosphate.,online information:Phosphoglycerate kinase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 2/5.,similarity:Belongs to the phosphoglycerate kinase family.,subunit:Monomer.,tissue specificity:Testis
	Background	arose via retrotransposition of the phosphoglycerate kinase 1 gene, and is expressed specifically in the testis. Initially assumed to be a pseudogene, the encoded protein is actually a functional phosphoglycerate kinase that catalyzes the reversible conversion of 1,3-bisphosphoglycerate to 3-phosphoglycerate, during the Embden-Meyerhof-Parnas pathway of glycolysis, in the later stages of



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PGK2 Monoclonal Antibody