



UB2L3 Monoclonal Antibody

Catalog No	YP-mAb-06363
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	UBE2L3 UBCE7 UBCH7
Protein Name	Ubiquitin-conjugating enzyme E2 L3 (EC 6.3.2.19) (L-UBC) (UbcH7) (Ubiquitin carrier protein L3) (Ubiquitin-conjugating enzyme E2-F1) (Ubiquitin-protein ligase L3)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	UB2L3 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	16kD
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Ubiquitous, with highest expression in testis.
Function	catalytic activity:ATP + ubiquitin + protein lysine = AMP + diphosphate + protein N-ubiquityllysine.,caution:PubMed:10760570 reported that UBE2L1, UBE2L2 and UBE2L4 are most likely pseudogenes and the only expressed member of this subfamily seems to be UBE2L3.,function:Catalyzes the covalent attachment of ubiquitin to other proteins. Mediates the selective degradation of short-lived and abnormal proteins. Functions in the E6/E6-AP-induced ubiquitination of p53/TP53.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the ubiquitin-conjugating enzyme family.,subunit:Bounds UBE3A, HEI10, CBL, ZAP70, RNF19A, RNF19B and RNF144B.,tissue specificity:Ubiquitous, with highest expression in testis.,
Background	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s). This



gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is demonstrated to participate in the ubiquitination of p53, c-Fos, and the NF- κ B precursor p105 in vitro. Several alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using UB2L3 Monoclonal Antibody