



MAGAB Monoclonal Antibody

Catalog No	YP-mAb-06470
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	MAGEA11 MAGE11
Protein Name	Melanoma-associated antigen 11 (Cancer/testis antigen 1.11) (CT1.11) (MAGE-11 antigen)
Immunogen	Synthesized peptide derived from human protein . at AA range: 30-110
Specificity	MAGAB Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	47kD
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Expressed in tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma. Expressed in testis, ovary, prostate, cancerous prostate, breast and adrenal tissue.
Function	function:Acts as androgen receptor co-regulator that increases androgen receptor activity by modulating the receptors interdomain interaction. May play a role in embryonal development and tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma. Expressed in testis, ovary, prostate, cancerous prostate, breast and adrenal tissue.,
Background	MAGE family member A11(MAGEA11) Homo sapiens This gene is a member of the MAGEA gene family. The members of this family encode proteins with 50 to 80% sequence identity to each other. The promoters and first exons of the MAGEA genes show considerable variability, suggesting that the existence of this gene family enables the same function to be expressed under different transcriptional controls. The MAGEA genes are clustered at chromosomal location Xq28. They have been implicated in some hereditary disorders, such as




dyskeratosis congenita. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using MAGAB Monoclonal Antibody