



# GAR1 Monoclonal Antibody

Catalog No	YP-mAb-06580
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GAR1 NOLA1
Protein Name	H/ACA ribonucleoprotein complex subunit 1 (Nucleolar protein family A member 1) (snoRNP protein GAR1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 110-190
Specificity	GAR1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	23kD
Cell Pathway	Nucleus, nucleolus. Nucleus, Cajal body. Also localized to Cajal bodies (coiled bodies).
Tissue Specificity	Placenta,
Function	domain:Interaction with SMN1 requires at least one of the RGG-box regions.,function:Required for ribosome biogenesis and telomere maintenance. Part of the H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which catalyzes pseudouridylation of rRNA. This involves the isomerization of uridine such that the ribose is subsequently attached to C5, instead of the normal N1. Each rRNA can contain up to 100 pseudouridine ("psi") residues, which may serve to stabilize the conformation of rRNAs. May also be required for correct processing or intranuclear trafficking of TERC, the RNA component of the telomerase reverse transcriptase (TERT) holoenzyme.,similarity:Belongs to the GAR1 family.,subcellular location:Also localized to Cajal bodies (coiled bodies),subunit:Part of the H/ACA small nucleolar ribonucleoprotein (H/ACA snoRNP) complex, which contains NHP2/NOLA2, GAR1/NOLA1, NOP10/N
Background	This gene is a member of the H/ACA snoRNPs (small nucleolar ribonucleoproteins) gene family. snoRNPs are involved in various aspects of rRNA processing and modification and have been classified into two families: C/D



and H/ACA. The H/ACA snoRNPs also include the DKC1, NOLA2 and NOLA3 proteins. These four H/ACA snoRNP proteins localize to the dense fibrillar components of nucleoli and to coiled (Cajal) bodies in the nucleus. Both 18S rRNA production and rRNA pseudouridylation are impaired if any one of the four proteins is depleted. These four H/ACA snoRNP proteins are also components of the telomerase complex. The encoded protein of this gene contains two glycine- and arginine-rich domains and is related to *Saccharomyces cerevisiae* Gar1p. Two splice variants have been found for this gene. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using GAR1 Monoclonal Antibody