




I20L2 Monoclonal Antibody

Catalog No	YP-mAb-06623
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	ISG20L2 HSD-38 HSD38
Protein Name	Interferon-stimulated 20 kDa exonuclease-like 2 (EC 3.1.-.-)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	I20L2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	38kD
Cell Pathway	Nucleus, nucleolus .
Tissue Specificity	Brain,Testis,
Function	function:3'-> 5'-exoribonuclease involved in ribosome biogenesis in the processing of the 12S pre-rRNA. Displays a strong specificity for a 3'-end containing a free hydroxyl group.,similarity:Contains 1 exonuclease domain.,
Background	This gene encodes a 3'-5' exoribonuclease that may be involved in the processing of the 12S pre-rRNA. Pseudogenes have been identified on chromosomes 6 and 11. [provided by RefSeq, Dec 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using I20L2 Monoclonal Antibody