






IN35 Monoclonal Antibody

| Catalog No | YP-mAb-06633 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | IFI35 IFP35 |
| Protein Name | Interferon-induced 35 kDa protein (IFP 35) (Ifi-35) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | IN35 Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 31kD |
| Cell Pathway | Cytoplasm . Nucleus . Secreted . Cytoplasmic IFI35 localizes in punctate granular structures (PubMed:10950963). Nuclear localization increased is stimulated by IFN-alpha (PubMed:8288566, PubMed:10950963). Extracelullar following secretion by macrophage (PubMed:29038465) |
| Tissue Specificity | Expressed in a wide range of cell types, including fibroblasts, macrophages, and epithelial cells. |
| Function | function:Not yet known.,induction:By interferon gamma.,similarity:Belongs to the NMI family.,subcellular location:Nuclear following IFN treatment.,subunit:Homodimer. Also interacts with B-ATF.,tissue specificity:In a wide range of cell types, including fibroblasts, macrophages, and epithelial cells., |
| Background | function:Not yet known.,induction:By interferon gamma.,similarity:Belongs to the NMI family.,subcellular location:Nuclear following IFN treatment.,subunit:Homodimer. Also interacts with B-ATF.,tissue specificity:In a wide range of cell types, including fibroblasts, macrophages, and epithelial cells., |
| matters needing attention | Avoid repeated freezing and thawing! |
| | |



UpingBio technology Co.,Ltd








Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using IN35 Monoclonal Antibody