




# XCL2 Monoclonal Antibody

<b>Catalog No</b>	YP-mAb-06662
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB
<b>Gene Name</b>	XCL2 SCYC2
<b>Protein Name</b>	Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein AA range: 1-50
<b>Specificity</b>	XCL2 Monoclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-1:2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	12kD
<b>Cell Pathway</b>	Secreted.
<b>Tissue Specificity</b>	Blood,Placenta,
<b>Function</b>	function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family.,
<b>Background</b>	function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western Blot analysis of various cells using XCL2  
Monoclonal Antibody