





C1QT5 Monoclonal Antibody

Catalog No	YP-mAb-07053
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	C1QTNF5 CTRP5 UNQ303/PRO344
Protein Name	Complement C1q tumor necrosis factor-related protein 5
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	C1QT5 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	26kD
Cell Pathway	Secreted .
Tissue Specificity	Brain,Fetal brain,Uterus,
Function	developmental stage:Expressed in fetal brain., disease:Defects in C1QTNF5 are a cause of late-onset retinal degeneration (LORD) [MIM:605670]. LORD is an autosomal dominant disorder characterized by onset in the fifth to sixth decade with night blindness and punctate yellow-white deposits in the retinal fundus, progressing to severe central and peripheral degeneration, with choroidal neovascularization and chorioretinal atrophy., disease:Defects in MFRP are the cause of microphthalmia MFRP-related (MCOPMFRP) [MIM:611040]. Microphthalmia is a clinically heterogeneous disorder of eye formation, ranging from small size of a single eye to complete bilateral absence of ocular tissues. Ocular abnormalities like opacities of the cornea and lens, scaring of the retina and choroid, cataract and other abnormalities like cataract may also be present. MCOPMFRP is characterized by posterior microphthalm
Background	This gene encodes a member of a family of proteins that function as components of basement membranes and may play a role in cell adhesion. Mutations in this gene have been associated with late-onset retinal degeneration. The protein may



UpingBio technology Co.,Ltd







be encoded by either a bicistronic transcript including sequence from the upstream membrane frizzled-related protein gene (MFRP), or by a monocistronic transcript expressed from an internal promoter. [provided by RefSeq, Jun 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using C1QT5 Monoclonal Antibody