



PON2 Monoclonal Antibody

Catalog No	YP-mAb-07061
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	PON2
Protein Name	Serum paraoxonase/arylesterase 2 (PON 2) (EC 3.1.1.2) (EC 3.1.1.81) (Aromatic esterase 2) (A-esterase 2) (Serum aryldialkylphosphatase 2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	PON2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	38kD
Cell Pathway	Membrane ; Peripheral membrane protein .
Tissue Specificity	Widely expressed with highest expression in liver, lung, placenta, testis and heart.
Function	catalytic activity:A phenyl acetate + H(2)O = a phenol + acetate.,catalytic activity:An aryl dialkyl phosphate + H(2)O = dialkyl phosphate + an aryl alcohol.,function:Hydrolyzes the toxic metabolites of a variety of organophosphorus insecticides. CaMABLE of hydrolyzing a broad spectrum of organophosphate substrates and a number of aromatic carboxylic acid esters (By similarity). Has antioxidant activity. Is not associated with high density lipoprotein. Prevents LDL lipid peroxidation, reverses the oxidation of mildly oxidized LDL, and inhibits the ability of MM-LDL to induce monocyte chemotaxis.,online information:The Singapore human mutation and polymorphism database,PTM:Glycosylated.,PTM:The signal sequence is not cleaved.,similarity:Belongs to the paraoxonase family.,tissue specificity:Widely expressed with highest expression in liver, lung, placenta, testis and heart.,
Background	This gene encodes a member of the paraoxonase gene family, which includes three known members located adjacent to each other on the long arm of chromosome 7. The encoded protein is ubiquitously expressed in human tissues,



membrane-bound, and may act as a cellular antioxidant, protecting cells from oxidative stress. Hydrolytic activity against acylhomoserine lactones, important bacterial quorum-sensing mediators, suggests the encoded protein may also play a role in defense responses to pathogenic bacteria. Mutations in this gene may be associated with vascular disease and a number of quantitative phenotypes related to diabetes. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PON2 Monoclonal Antibody