




LECT2 Monoclonal Antibody

Catalog No	YP-mAb-07066
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	LECT2
Protein Name	Leukocyte cell-derived chemotaxin-2 (LECT-2) (hLECT2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	LECT2 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	16kD
Cell Pathway	Cytoplasm . Secreted .
Tissue Specificity	Highly expressed in adult and fetal liver and weakly in testis. Not expressed in bone marrow.
Function	function:Has a neutrophil chemotactic activity. Also a positive regulator of chondrocyte proliferation.,induction:By phytohemagglutinin (PHA).,similarity:Belongs to the LECT2/MIM-1 family.,tissue specificity:Highly expressed in adult and fetal liver and weakly in testis. Not expressed in bone marrow.,
Background	This gene encodes a secreted, 16 kDa protein that acts as a chemotactic factor to neutrophils and stimulates the growth of chondrocytes and osteoblasts. This protein has high sequence similarity to the chondromodulin repeat regions of the chicken myb-induced myeloid 1 protein. A polymorphism in this gene may be associated with rheumatoid arthritis. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using LECT2 Monoclonal Antibody