





RAP1B Monoclonal Antibody

Catalog No YP-mAb-07099 Isotype IgG Reactivity Human;Rat;Mouse Applications WB Gene Name RAP1B OK/SW-cl.11 Protein Name Ras-related protein Rap-1b (GTP-binding protein smg p21B) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol. Cell junction. May shuttle between plasma membrane and cytosol, (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala, Brain, Chondrosarcoma, Colon adenocarcinoma, Placenta, Platele Function enzyme regulatio		
Reactivity Human;Rat;Mouse Applications WB Gene Name RAP1B OK/SW-cl.11 Protein Name Ras-related protein Rap-1b (GTP-binding protein smg p21B) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala, Brain, Chondrosarcoma, Colon adenocarcinoma, Placenta, Platele Function enzyme regulation: Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner . isamilarity. Belongs to the small GTP-ase superfamily. Members of this family regulate multiple cellular processes including oral advanced in a grower and provided in the grain of the plasma membrane and cytosol, subunit Interacts with SGSM1, SGSM2 and SGSM3. Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including oral advanced in regulating outside-in spanning in plate act ell signaling in plateits, splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9, [provided by RefSeq, Oct 2011],	Catalog No	YP-mAb-07099
Applications WB Gene Name RAP1B OK/SW-cl.11 Protein Name Ras-related protein Rap-1b (GTP-binding protein smg p21B) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity. Belongs to the small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including superfamily. Members of this family regulate multiple cellular processes including reportance and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating integrin-mediated cell signaling. This protein also plays a role in regulating unitspread integrin-mediated cell signaling. This protein also plays a role in regulating unitspread integrin-mediated cell signaling. This protein also plays a role in regulating unitspread integrin-mediated cell signaling. This protein also plays a role in regulating integrin-mediated cell signaling. This protein incedires. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9, [provided by RefSeq, Oct 2011].	Isotype	IgG
Gene Name RAP1B OK/SW-cl.11 Protein Name Ras-related protein Rap-1b (GTP-binding protein smg p21B) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH6 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma, Colon adenocarcinoma, Placenta, Platele Function enzyme regulation: Activated by guanine nucleotide-exchange small GTP-be small GTP-se superfamily. Ras family, subcellular location: May shuttle between plasma membrane and cytosol., subunit: Interacts with SGSM1, SGSM2 and SGSM3. Background This gene en	Reactivity	Human;Rat;Mouse
Protein Name Ras-related protein Rap-1b (GTP-binding protein smg p21B) Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH6 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala, Brain, Chondrosarcoma, Colon adenocarcinoma, Placenta, Platele enzyme regulation: Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner, similarity. Belongs to the small GTPase superfamily. Ras family, subcellular location: May shuttle between plasma membrane and cytosol. subunit: Interacts with SGSM1, SGSM2 and SGSM3. Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell alchesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein in also plays a role in regulating outside-in signaling localizes to cellular processes including cell alchesion and growth and differentiation. This protein localizes to cellular processes including on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011],	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 110-190 Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma, Colon adenocarioma, Placenta, Platele Function enzyme regulation-Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner, similarity:Belongs to the small GTPase superfamily. Ras family, subcellular location:May shuttle between plasma membrane and cytosol., subunit:Interacts with SGSM1, SGSM2 and SGSM3. Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including newbrones and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in reg	Gene Name	RAP1B OK/SW-cl.11
Specificity RAP1B Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner. similarity: Belongs to the small GTPase superfamily. Ras family, subcellular location:May shuttle between plasma membrane and cytosol, subunit:Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9, [provided by RefSeq, Oct 2011],	Protein Name	Ras-related protein Rap-1b (GTP-binding protein smg p21B)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed: 3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala, Brain, Chondrosarcoma, Colon adenocarcinoma, Placenta, Platele Function enzyme regulation: Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner, similarity: Belongs to the small GTPase superfamily. Ras family, subcellular location: May shuttle between plasma membrane and cytosol, subunit: Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011],	Immunogen	Synthesized peptide derived from human protein . at AA range: 110-190
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:31414412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120) . Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a CAMP-dependent manner., similarity:Belongs to the small GTPase superfamily. Ras family, subcellular location:May shuttle between plasma membrane and cytosol., subunit:Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-pinding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011]. <tr< th=""><th>Specificity</th><th>RAP1B Monoclonal Antibody detects endogenous levels of protein.</th></tr<>	Specificity	RAP1B Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcioma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol, subunit:Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011],	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala, Brain, Chondrosarcoma, Colon adenocarcinoma, Placenta, Platele Function enzyme regulation: Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner. similarity: Belongs to the small GTPase superfamily. Ras family. subcellular location: May shuttle between plasma membrane and cytosol., subunit: Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], Matters needing Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120) . Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family, subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1 SGSM2 and SGSM3. Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], matters needing Avoid repeated freezing and thawing!	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 20kD Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-1:2000
Storage Stability -20°C/1 year Observed Band Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120) Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family, subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3, Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Observed Band Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3., This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane . Cytoplasm, cytosol . Cell junction . May shuttle between plasma membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120) Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3., This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], Matters needing Avoid repeated freezing and thawing!	Synonyms	
membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is required for its localization to the cell junction (PubMed:20332120). Tissue Specificity Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3., This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], Matters needing Avoid repeated freezing and thawing!	Observed Band	20kD
Function enzyme regulation:Activated by guanine nucleotide-exchange factor (GEF) EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma membrane and cytosol.,subunit:Interacts with SGSM1, SGSM2 and SGSM3., This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], Matters needing Avoid repeated freezing and thawing!	Cell Pathway	membrane and cytosol (PubMed:3141412). Presence of KRIT1 and CDH5 is
EPÁC2 in a cAMP-dependent manner., similarity:Belongs to the small GTPase superfamily. Ras family., subcellular location:May shuttle between plasma membrane and cytosol., subunit:Interacts with SGSM1, SGSM2 and SGSM3., Background This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], Matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Amygdala,Brain,Chondrosarcoma,Colon adenocarcinoma,Placenta,Platele
superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011], matters needing Avoid repeated freezing and thawing!	Function	EPAC2 in a cAMP-dependent manner.,similarity:Belongs to the small GTPase superfamily. Ras family.,subcellular location:May shuttle between plasma
	Background	superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found
		Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd







Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



