








DKKL1 Monoclonal Antibody

Catalog No	YP-mAb-07109
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DKKL1 SGY1 UNQ735/PRO1429
Protein Name	Dickkopf-like protein 1 (Cancer/testis antigen 34) (CT34) (Protein soggy-1) (SGY-1)
Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Specificity	DKKL1 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Descritor	>00%
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 26kD Secreted . Cytoplasmic vesicle, secretory vesicle, acrosome . Localized
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 26kD Secreted . Cytoplasmic vesicle, secretory vesicle, acrosome . Localized specifically to the crescent shaped acrosome at the apex of the sperm head More highly expressed in adult testis than in fetal testis. Exclusively expressed in the testis (at protein level). Intense expression in stages II, III and IV of
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 26kD Secreted . Cytoplasmic vesicle, secretory vesicle, acrosome . Localized specifically to the crescent shaped acrosome at the apex of the sperm head More highly expressed in adult testis than in fetal testis. Exclusively expressed in the testis (at protein level). Intense expression in stages II, III and IV of spermatogenesis, whereas expression is lower in stage I. PTM:N-glycosylated.,similarity:To the N-terminal section of DKK-3., The dickkopf protein family interacts with the Wnt signaling pathway and its members are characterized by two conserved cysteine-rich domains. This gene encodes a secreted protein that has low sequence similarity to the dickkopf-3
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	26kD Secreted . Cytoplasmic vesicle, secretory vesicle, acrosome . Localized specifically to the crescent shaped acrosome at the apex of the sperm head More highly expressed in adult testis than in fetal testis. Exclusively expressed in the testis (at protein level). Intense expression in stages II, III and IV of spermatogenesis, whereas expression is lower in stage I. PTM:N-glycosylated.,similarity:To the N-terminal section of DKK-3., The dickkopf protein family interacts with the Wnt signaling pathway and its members are characterized by two conserved cysteine-rich domains. This gene encodes a secreted protein that has low sequence similarity to the dickkopf-3 protein. Multiple alternatively spliced transcript variants encoding distinct isoforms
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background	26kD Secreted . Cytoplasmic vesicle, secretory vesicle, acrosome . Localized specifically to the crescent shaped acrosome at the apex of the sperm head More highly expressed in adult testis than in fetal testis. Exclusively expressed in the testis (at protein level). Intense expression in stages II, III and IV of spermatogenesis, whereas expression is lower in stage I. PTM:N-glycosylated.,similarity:To the N-terminal section of DKK-3., The dickkopf protein family interacts with the Wnt signaling pathway and its members are characterized by two conserved cysteine-rich domains. This gene encodes a secreted protein that has low sequence similarity to the dickkopf-3 protein. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Oct 2010],





Products Images



Western Blot analysis of various cells using DKKL1 Monoclonal Antibody