




HRK Monoclonal Antibody

Catalog No	YP-mAb-07120
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	HRK BID3
Protein Name	Activator of apoptosis harakiri (BH3-interacting domain-containing protein 3) (Neuronal death protein DP5)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 1-50
Specificity	HRK Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	10kD
Cell Pathway	Membrane; Single-pass membrane protein. Mitochondrion.
Tissue Specificity	
Function	domain:The BH3 motif is required for the induction of cell death.,function:Activates apoptosis and interacts selectively with survival-promoting proteins Bcl-2 and Bcl-X(L).,
Background	harakiri, BCL2 interacting protein(HRK) Homo sapiens This gene encodes a member of the BCL-2 protein family. Members of this family are involved in activating or inhibiting apoptosis. The encoded protein localizes to intracellular membranes. This protein promotes apoptosis by interacting with the apoptotic inhibitors BCL-2 and BCL-X(L) via its BH3 domain. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Oct 2012],
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using HRK Monoclonal Antibody