





## **ELAF Monoclonal Antibody**

Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  Function (domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase		
Reactivity Human;Rat:Mouse;  Applications WB  Gene Name PI3 WAP3 WFDC14  Protein Name Elafin (Elastase-specific inhibitor) (ESI) (Peptidase inhibitor 3) (PI-3) (Protease inhibitor WAP3) (Skin-derived antifleukoproteinase) (SKALP) (WAP four-disulfide core domain protein 14)  Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent leaflates-mediated tissue proteolysis, similarity:Contains 1 WAP domain, and st thus a member of the WFDC domain family, Most WFDC gene members are localized to chromosome 20412-413 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial [Inpopolysaccharides and cytokines.] provided by gene is upgulated by bacterial [Inpopolysaccharides and cytokines.]	Catalog No	YP-mAb-07235
Applications WB  Gene Name PI3 WAP3 WFDC14  Protein Name Elafin (Elastase-specific inhibitor) (ESI) (Peptidase inhibitor 3) (PI-3) (Protease inhibitor WAP3) (Skin-derived antileukoproteinase) (SKALP) (WAP four-disulfide core domain protein 14)  Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain retracellular matrix proteins, function Neutrophil and pancreafic elastase-specific inhibitor of skin. It may prevent leaflafic ovalently to specific site on extracellular matrix proteins, function. Neutrophil and pancreafic elastase-specific inhibitor of skin. It may prevent leaflase-mediated tissue proteolysis, similarity. Contains 1 WAP domain.  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-distilfier or (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20112-413 in two clusters centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial (Inpo)polysaccharides and cytokines.	Isotype	IgG
Gene Name PI3 WAP3 WFDC14  Protein Name Elafin (Elastase-specific inhibitor) (ESI) (Peptidase inhibitor 3) (PI-3) (Protease inhibitor WAP3) (Skin-derived antileukoproteinase) (SKALP) (WAP four-disulfide core domain protein 14)  Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum, domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain received and substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function: Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis, similarity: Contains 1 WAP domain.  Background This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines, provided by	Reactivity	Human;Rat;Mouse;
Protein Name  Elafin (Elastase-specific inhibitor) (ESI) (Peptidase inhibitor 3) (PI-3) (Protease inhibitor WAP3) (Skin-derived antileukoproteinase) (SKALP) (WAP four-disulfide core domain protein 14)  Immunogen  Synthesized peptide derived from human protein . at AA range: 40-120  Specificity  ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Monoclonal, Mouse, IgG  Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  12kD  Cell Pathway  Secreted.  Tissue Specificity  Brain, Keratinocyte, Placenta, Sputum, Stratum corneum, domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain recention moiety) and the elastase inhibitor domain. The transglutaminase substrate domain substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function: Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis, similarity: Contains 1 WAP domain.  Background  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disusfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is uppelated by bacterial lipopolysaccharides and cytokines, provided by	Applications	WB
core domain profein 14)  Immunogen Synthesized peptide derived from human protein . at AA range: 40-120  Specificity ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain matrix proteins, function: Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis, similarity: Contains 1 WAP domain.  Background This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20(12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric and cytokines. [provided by secretal lippoplysaccharides and cytokines. [provided by gene is upgulated by bacterial lippoplysaccharides and cytokines. [provided by gene is upgulated by bacterial lippoplysaccharides and cytokines. [provided by gene is upgulated by bacterial lippoplysaccharides and cytokines. [provided by gene is upgulated by bacterial lippoplysaccharides and cytokines. [provided by	Gene Name	PI3 WAP3 WFDC14
Specificity ELAF Monoclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  function domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function: Neutrophil and pancreatic elastase-specific inhibitor of sin. It may prevent elastase-mediated tissue proteolysis, similarity: Contains 1 WAP domain.  Background This gene encodes an elastase-specific inhibitor that functions as an antimitrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by bacterial lipopolysaccharides and cytokines. [provided by bacterial lipopolysaccharides and cytokines. [provided by sacterial lip	Protein Name	Elafin (Elastase-specific inhibitor) (ESI) (Peptidase inhibitor 3) (PI-3) (Protease inhibitor WAP3) (Skin-derived antileukoproteinase) (SKALP) (WAP four-disulfide core domain protein 14)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  Function domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis, similarity:Contains 1 WAP domain.  Background  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lippoplysaccharides and cytokines. [provided by	Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Source         Monoclonal, Mouse,IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band         12kD           Cell Pathway         Secreted.           Tissue Specificity         Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,           domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalentity to specific site on extracellular matrix proteins, function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis. similarity:Contains 1 WAP domain.           Background         This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Specificity	ELAF Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000 Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms Observed Band 12kD Cell Pathway Secreted.  Tissue Specificity Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain roveracelic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis, similarity:Contains 1 WAP domain.  Background This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokiness. [provided by	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis., similarity:Contains 1 WAP domain.  Background This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       12kD         Cell Pathway       Secreted.         Tissue Specificity       Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,         Function       domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins, function: Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis, similarity: Contains 1 WAP domain.         Background       This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  function domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain reverse as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins. function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  Background This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Dilution	WB 1:500-1:2000
Storage Stability  -20°C/1 year  Synonyms  Observed Band  12kD  Cell Pathway  Secreted.  Tissue Specificity  Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific site on extracellular matrix proteins, function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  Background  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Concentration	1 mg/ml
Synonyms  Observed Band  12kD  Cell Pathway  Secreted.  Tissue Specificity  Brain, Keratinocyte, Placenta, Sputum, Stratum corneum,  function  domain: Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins., function: Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis., similarity: Contains 1 WAP domain.,  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Purity	≥90%
Cell Pathway  Secreted.  Tissue Specificity  Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins.,function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Storage Stability	-20°C/1 year
Cell Pathway  Secreted.  Tissue Specificity  Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins.,function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  Background  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Synonyms	
Tissue Specificity  Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,  domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins.,function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Observed Band	12kD
domain:Consists of two domains: the transglutaminase substrate domain (cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins.,function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Cell Pathway	Secreted.
(cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins.,function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue proteolysis.,similarity:Contains 1 WAP domain.,  This gene encodes an elastase-specific inhibitor that functions as an antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Tissue Specificity	Brain,Keratinocyte,Placenta,Sputum,Stratum corneum,
antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by	Function	(cementoin moiety) and the elastase inhibitor domain. The transglutaminase substrate domain serves as an anchor to localize elafin covalently to specific sites on extracellular matrix proteins.,function:Neutrophil and pancreatic elastase-specific inhibitor of skin. It may prevent elastase-mediated tissue
	Background	antimicrobial peptide against Gram-positive and Gram-negative bacteria, and fungal pathogens. The protein contains a WAP-type four-disulfide core (WFDC) domain, and is thus a member of the WFDC domain family. Most WFDC gene members are localized to chromosome 20q12-q13 in two clusters: centromeric and telomeric. This gene belongs to the centromeric cluster. Expression of this gene is upgulated by bacterial lipopolysaccharides and cytokines. [provided by



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using ELAF Monoclonal Antibody