



FABP5 Monoclonal Antibody

Catalog No	YP-mAb-07244
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB
Gene Name	FABP5
Protein Name	Fatty acid-binding protein, epidermal (Epidermal-type fatty acid-binding protein) (E-FABP) (Fatty acid-binding protein 5) (Psoriasis-associated fatty acid-binding protein homolog) (PA-FABP)
Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Specificity	FABP5 Monoclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	14kD
Cell Pathway	Cytoplasm . Nucleus . Cell junction, synapse . Cell junction, synapse, postsynaptic density . Secreted . Localizes primarily to the cytoplasm. Upon certain ligand binding, a conformation change exposes a nuclear localization motif and the protein is transported into nucleus (PubMed:24692551). Secreted by astrocytes, but not by neurons (By similarity). .
Tissue Specificity	Keratinocytes; highly expressed in psoriatic skin (PubMed:8092987). Expressed in brain gray matter (PubMed:21395585).
Function	domain:Forms a beta-barrel structure that accommodates the hydrophobic ligand in its interior.,function:High specificity for fatty acids. Highest affinity for C18 chain length. Decreasing the chain length or introducing double bonds reduces the affinity. May be involved in keratinocyte differentiation.,similarity:Belongs to the calycin superfamily. Fatty-acid binding protein (FABP) family.,tissue specificity:Keratinocytes; highly expressed in psoriatic skin.,
Background	This gene encodes the fatty acid binding protein found in epidermal cells, and was first identified as being upregulated in psoriasis tissue. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. FABPs may play roles in



fatty acid uptake, transport, and metabolism. Polymorphisms in this gene are associated with type 2 diabetes. The human genome contains many pseudogenes similar to this locus.[provided by RefSeq, Feb 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using FABP5 Monoclonal Antibody