





GPR84 Monoclonal Antibody

kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1		
Reactivity Human;Rat;Mouse; Applications WB Gene Name GPR84 EX33 Protein Name G-protein coupled receptor 84 (Inflammation-related G-protein coupled receptor EX33) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity GPR84 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, and essingohlis relative to T- or B-lymphocytes. Function function: Orphan receptor, similarity Belongs to the G-protein coupled receptor 1 family, tissue specificity, Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Background function: Orphan receptor, similarity-Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes.	Catalog No	YP-mAb-07391
Applications Gene Name GPR84 EX33 Protein Name G-protein coupled receptor 84 (Inflammation-related G-protein coupled receptor EX33) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity GPR84 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kiney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function: Orphan receptor, similarity, Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Background	Isotype	IgG
Gene Name GPR84 EX33 Protein Name G-protein coupled receptor 84 (Inflammation-related G-protein coupled receptor EX33) Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity GPR84 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kiney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophilis and eosinophilis relative to T- or B-lymphocytes. Function function:Orphan receptor, similarity.Belongs to the G-protein coupled receptor 1 family, tissue specificity; Expressed in brain, heart, muscle, colon, thymus, spleen,	Reactivity	Human;Rat;Mouse;
Protein Name G-protein coupled receptor 84 (Inflammation-related G-protein coupled receptor EX33)	Applications	WB
Immunogen Synthesized peptide derived from human protein . at AA range: 230-310 Specificity GPR84 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophilis and eosinophilis relative to T- or B-lymphocytes. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophilis and eosinophilis, in comparison with that in T- or B-lymphocytes. Background function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophilis and eosinophilis, in comparison with that in T- or B-lymphocytes.	Gene Name	GPR84 EX33
Specificity GPR84 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Background function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils in comparison with that in T- or B-lymphocytes.	Protein Name	G-protein coupled receptor 84 (Inflammation-related G-protein coupled receptor EX33)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Background function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils in comparison with that in T- or B-lymphocytes.	Immunogen	Synthesized peptide derived from human protein . at AA range: 230-310
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-Iymphocytes. Function function:Orphan receptor, similarity:Belongs to the G-protein coupled receptor 1 family, tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Background function:Orphan receptor, similarity:Belongs to the G-protein coupled receptor 1 family, tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils <td>Specificity</td> <td>GPR84 Monoclonal Antibody detects endogenous levels of protein.</td>	Specificity	GPR84 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor, similarity:Belongs to the G-protein coupled receptor 1 family, lissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Function:Orphan receptor, similarity:Belongs to the G-protein coupled receptor 1 family, lissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophilis	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor_similarity:Belongs to the G-protein coupled receptor 1 family_tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes. Background function:Orphan receptor_similarity:Belongs to the G-protein coupled receptor 1 family_tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophilis niceptor. Similarity:Belongs to the G-protein coupled receptor 1 family_tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophilis niceptor. Similarity:Belongs to the G-protein coupled receptor 1 family_tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophilis niceptor. In the time the protein coupled receptor in neutrophilis niceptor and the protein cou	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function: Orphan receptor., similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes.	Purification	· · · · · · · · · · · · · · · · · · ·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils in the protein coupled receptor 1 family, tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Dilution	WB 1:500-1:2000
Synonyms Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Concentration	1 mg/ml
Observed Band 43kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function: Orphan receptor., similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function: Orphan receptor., similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Purity	≥90%
Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Synonyms	
Tissue Specificity Expressed predominantly in hematopoietic tissues. High levels detected in the bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Observed Band	43kD
bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils and eosinophils relative to T- or B-lymphocytes. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Cell Pathway	Cell membrane; Multi-pass membrane protein.
family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils and eosinophils, in comparison with that in T- or B-lymphocytes., Background function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Tissue Specificity	bone marrow and lower levels in the peripheral leukocytes and lung. Also expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta and intestine. Within the leukocyte population expression is higher in neutrophils
family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils	Function	family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils
	Background	family.,tissue specificity:Expressed in brain, heart, muscle, colon, thymus, spleen, kidney, liver, placenta, intestine, bone marrow, lung and peripheral blood leukocytes. Within leukocyte population more abundant expression in neutrophils



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GPR84 Monoclonal Antibody