

OR1J4 Monoclonal Antibody

Catalog No YP-mAb-07452 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name OR1J4 Protein Name Olfactory receptor 1J4 (HTPCRX01) (Olfactory receptor OR9-21) Immunogen Synthesized peptide derived from part region of human protein Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptor family 1 subfamily J member 4(OR1,J4) Homo sapiens Olfactory receptor geno		
Reactivity Human;Rat;Mouse; Applications WB Gene Name OR1J4 Protein Name Olfactory receptor 1J4 (HTPCRX01) (Olfactory receptor OR9-21) Immunogen Synthesized peptide derived from part region of human protein Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-romatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones,Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens olfactory receptor a large family of G-protein roupled receptor 1 response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein coupled receptors or receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many heurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms is independent of other organisms. [provided by the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms is independent of other organisms. [provided by	Catalog No	YP-mAb-07452
Applications WB Gene Name OR1J4 Protein Name Olfactory receptor 1J4 (HTPCRX01) (Olfactory receptor OR9-21) Immunogen Synthesized peptide derived from part region of human protein Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptor sinteract with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor septor are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors of this organism is independent of other organisms. [provided by	Isotype	IgG
Gene Name OR1J4 Protein Name Olfactory receptor 1J4 (HTPCRX01) (Olfactory receptor OR9-21) Immunogen Synthesized peptide derived from part region of human protein Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function: Odorant receptor ,,similarity: Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms, provided by	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 1J4 (HTPCRX01) (Olfactory receptor OR9-21) Immunogen Synthesized peptide derived from part region of human protein Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors share a return of protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a return of proteins of the organism is indeponent of other organisms, provided by	Applications	WB
Immunogen Synthesized peptide derived from part region of human protein Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Function function:Odorant receptor.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1,J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organisms is independent of o	Gene Name	OR1J4
Specificity OR1J4 Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that tringgers the perception of a semil. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor gene family is the largest in the geno	Protein Name	Olfactory receptor 1J4 (HTPCRX01) (Olfactory receptor OR9-21)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms in independent of other organisms. [provided by	Immunogen	Synthesized peptide derived from part region of human protein
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Specificity	OR1J4 Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms, [provided by	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors warising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Source	Monoclonal, Mouse,lgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function: Odorant receptor,similarity: Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Dilution	WB 1:500-1:2000
Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Concentration	1 mg/ml
Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purity	≥90%
Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity PCR rescued clones, Testis, Function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. PCR rescued clones, Testis, Function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Synonyms	
Tissue Specificity PCR rescued clones, Testis, function function:Odorant receptor ., similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Observed Band	34kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background Olfactory receptor family 1 subfamily J member 4(OR1J4) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Tissue Specificity	PCR rescued clones, Testis,
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
	Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using OR1J4 Monoclonal Antibody