

## P2RY1 Monoclonal Antibody

| Catalog No         | YP-mAb-07527   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Rat;Mouse  |
| Applications       | WB   |
| Gene Name          | P2RY1  |
| Protein Name       | P2Y purinoceptor 1 (P2Y1) (ATP receptor) (Purinergic receptor)   |
| Immunogen          | Synthesized peptide derived from human protein . at AA range: 130-210  |
| Specificity        | P2RY1 Monoclonal Antibody detects endogenous levels of protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| Source             | Monoclonal, Mouse,IgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-1:2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           |  |
| Observed Band      | 41kD   |
| Cell Pathway       | Cell membrane ; Multi-pass membrane protein .  |
| Tissue Specificity | Blood,Brain,Lung,Placenta,Platelet,  |
| Function           | function:Receptor for extracellular adenine nucleotides such as ATP and ADP. In platelets binding to ADP leads to mobilization of intracellular calcium ions via activation of phospholipase C, a change in platelet shape, and probably to platelet aggregation.,induction:Repressed by the P2Y1 receptor-specific antagonists A3P5PS, A3P5P and A2P5P. These inhibit calcium ion mobilization and shape change in platelets.,similarity:Belongs to the G-protein coupled receptor 1 family.,   |
| Background         | purinergic receptor P2Y1(P2RY1) Homo sapiens The product of this gene belongs to the family of G-protein coupled receptors. This family has several receptor subtypes with different pharmacological selectivity, which overlaps in some cases, for various adenosine and uridine nucleotides. This receptor functions as a receptor for extracellular ATP and ADP. In platelets binding to ADP leads to mobilization of intracellular calcium ions via activation of phospholipase C, a change in platelet shape, and probably to platelet aggregation. [provided by RefSeq, Jul 2008], |



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using P2RY1 Monoclonal Antibody