

## **OR7A2 Monoclonal Antibody**

YP-mAb-07626
IgG
Human;Rat;Mouse;
WB
OR7A2P OR7A2 OR7A7
Putative olfactory receptor 7A2 (Putative olfactory receptor 7A7)
Synthesized peptide derived from human protein . at AA range: 170-250
OR7A2 Monoclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
34kD
Cell membrane; Multi-pass membrane protein.
caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
caution:Could be the product of a pseudogene.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,  Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],



## UpingBio technology Co.,Ltd








**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using OR7A2 Monoclonal Antibody