





PCY1A Monoclonal Antibody

| Catalog No | YP-mAb-07755 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | PCYT1A CTPCT PCYT1 |
| Protein Name | Choline-phosphate cytidylyltransferase A (EC 2.7.7.15) (CCT-alpha) (CTP:phosphocholine cytidylyltransferase A) (CCT A) (CT A) (Phosphorylcholine transferase A) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | PCY1A Monoclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 40kD |
| Cell Pathway | Cytoplasm, cytosol . Membrane ; Peripheral membrane protein . Endoplasmic reticulum . Nucleus . It can interconvert between an inactive cytosolic form and an active membrane-bound form |
| Tissue Specificity | Brain, placenta, liver, fetal and adult lung. |
| Function | catalytic activity:CTP + choline phosphate = diphosphate + CDP-choline.,enzyme regulation:By phosphorylation.,function:Controls phosphatidylcholine synthesis.,pathway:Phospholipid metabolism; phosphatidylcholine biosynthesis; phosphatidylcholine from phosphocholine: step 1/2.,PTM:The serine residues of the C-terminus are phosphorylated. The inactive soluble form is stabilized by phosphorylation, the active membrane bound form is promoted by anionic lipids or diacylglycerol, and is stabilized by dephosphorylation.,similarity:Belongs to the cytidylyltransferase family.,subcellular location:It can interconvert between an inactive cytosolic form and an active membrane-bound form.,subunit:Homodimer; disulfide-linked., |
| Background | This gene belongs to the cytidylyltransferase family and is involved in the regulation of phosphatidylcholine biosynthesis. Mutations in this gene are associated with spondylometaphyseal dysplasia with cone-rod dystrophy. |



UpingBio technology Co.,Ltd







Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Aug 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PCY1A Monoclonal Antibody