





PODO Monoclonal Antibody

Catalog No YP-mAb-07756 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name NPHS2 Protein Name Podocin Immunogen Synthesized peptide derived from part region of human protein Specificity PODO Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane ; Peripheral membrane protein ; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and live years, resistance		
Reactivity	Catalog No	YP-mAb-07756
Applications WB Gene Name NPHS2 Protein Name Podocin Immunogen Synthesized peptide derived from part region of human protein Specificity PODO Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. but the protein specific processive steroid-resistant nephrotic syndrome (SRN) [MIM:600995], SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomeruloselorosis is permeability, acting projects by a silver between the plasma membrane and the cytoskeleton, similarity Belongs to the band 7/mec-2 family, subunit Interacts with nephrin/NPI-61 and KIRREL. Interacts directly with CD2AP. Interacts with nephrin/NPI-61 and KIRREL. Interacts directly with CD2AP interacts with nephrin/NPI-61 and KIRREL interacts directly variants. [provider syndrome, Alternative splicing results in multiple transcript variants. [provide by RefSeq. Jul Alternative splicing results in multiple transcript variants. [provided by RefSeq. Jul Alternative splicing results in multiple transcript variants. [provided by RefSeq. Jul Alternative splicing results in multiple transcript variants. [provided by RefSeq. Jul Alternative splicing results in multiple transcript variants. [provided by RefSeq. Jul Alternative splicing results in multiple transcript variants. [provided by RefSeq. Jul	Isotype	IgG
Protein Name	Reactivity	Human;Rat;Mouse;
Immunogen Synthesized peptide derived from part region of human protein	Applications	WB
Immunogen Synthesized peptide derived from part region of human protein Specificity PODO Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Cell Pathway [isoform 1]: Cell membrane ; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum . Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and live years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulas permeability, acting probably as a linker between the plasm amembrane and the cytoskeletonsimilarity.Belongs to the band 7/mec-2 family, subunit.Interacts with DDN, tissue specificity-Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruliar permeability. Acting probably as a linker between the plasm amembrane and mature kidney glo	Gene Name	NPHS2
Specificity PODO Monoclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease: Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages, function: Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton., similarity Belongs to the band 7/mec-2 family, subunit: Interacts with DDN., tissue specificity: Almost exclusively expressed in the podocytes of fetal and mature kidney glomerulis. This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome results in multiple transcript variants. [provided by RefSeq, Jul Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Protein Name	Podocin
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease: Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages. function: Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton., similarity, Belongs to the band 7/mec-2 family, subunit: Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN., itssue specificity: Almost exclusively expressed in the podocytes of fetal and mature kidney glomerulii. Background This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrolic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Immunogen	Synthesized peptide derived from part region of human protein
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIMi:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages, function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton. similarity: Belongs to the band 7/mez-2 family, subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Specificity	PODO Monoclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages. function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton. similarity:Belongs to the band 7/me-2-2 family, subunit:Interacts with DDN., tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Background This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease: Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages, function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton, similarity. Belongs to the band 7/mec-2 family, subunit: Interacts with DDN, tissue specificity: Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Background This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton, similarity:Belongs to the band 7/mec-2 family, subunit: Interacts with nephrin/NPHS1 and KIRREL. Interacts directly the CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages, function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton. similarity:Belongs to the band 7/mec-2 family., subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN., tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomerulli. Background This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Dilution	WB 1:500-1:2000
Synonyms Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein.; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages., function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton., similarity:Belongs to the band 7/mec-2 family., subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN., tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Background This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Concentration	1 mg/ml
Observed Band 42kD Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein.; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli., This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Purity	≥90%
Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein.; [Isoform 2]: Endoplasmic reticulum. Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli., This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Storage Stability	-20°C/1 year
Cell Pathway [Isoform 1]: Cell membrane; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum. Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli., This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Synonyms	
Tissue Specificity Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli. Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli., This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Observed Band	42kD
Function disease:Defects in NPHS2 are the cause of autosomal recessive steroid-resistant nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli., This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Cell Pathway	[Isoform 1]: Cell membrane ; Peripheral membrane protein .; [Isoform 2]: Endoplasmic reticulum .
nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and mature kidney glomeruli., Background This gene encodes a protein that plays a role in the regulation of glomerular permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Tissue Specificity	
permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul	Function	nephrotic syndrome (SRN) [MIM:600995]. SRN is characterized by onset between three months and five years, resistance to steroid therapy and rapid progression to end-stage renal disease. Focal segmental glomerulosclerosis is present at later stages.,function:Plays a role in the regulation of glomerular permeability, acting probably as a linker between the plasma membrane and the cytoskeleton.,similarity:Belongs to the band 7/mec-2 family.,subunit:Interacts with nephrin/NPHS1 and KIRREL. Interacts directly with CD2AP. Interacts with DDN.,tissue specificity:Almost exclusively expressed in the podocytes of fetal and
	Background	permeability. Mutations in this gene cause steroid-resistant nephrotic syndrome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PODO Monoclonal Antibody