


NPRL3 mouse mAb

nan NPRL3 AA range: 461-511 rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide.
rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide. In mouse antiserum by
rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide. In mouse antiserum by
rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide. In mouse antiserum by
rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide. In mouse antiserum by
rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide. In mouse antiserum by
rels of NPRL3 at Human/Mouse I, 0.5% BSA and 0.170% sodium azide. In mouse antiserum by
I, 0.5% BSA and 0.170% sodium azide.
n mouse antiserum by
•
•
-specific immunogen.
otein (-14 gene protein) (Alpha-globin rotein) (Protein CGTHBA)
lysosomes is amino acid-independent
rontal lobe cortex as well as in the temporal, 27173016, PubMed:26505888).
an genome assembly but is present in all Belongs to the UPF0171 family.,tissue
not known. [provided by RefSeq, Aug
ogical reaction related experiments. For nical personnel.



Products Images



Western Blot analysis of various cells using NPRL3 mouse mAb