





## INT4 mouse mAb

Catalog No         YP-mAb-08147           Isotype         IgG           Reactivity         Human; Mouse           Applications         WB           Gene Name         INTS4 MSTP093           Protein Name         INT4           Immunogen         Synthesized peptide derived from human INT4 AA range: 10-60           Specificity         This antibody detects endogenous levels of INT4 at Human/Mouse           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.           Source         Monocional, Mouse IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Integrator complex subunit 4 (Int4)           Observed Band         105kD           Cell Pathway         Nucleus .           Tissue Specificity         Aorta, Brain, Muscle, Ovary, Uterus,           Function         function: Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3-box-dependent processing. The Integrator complex is associated with the C-terminal domain class and processin		
Reactivity Human; Mouse  Applications WB  Gene Name INTS4 MSTP093  Protein Name INT4  Immunogen Synthesized peptide derived from human INT4 AA range: 10-60  Specificity This antibody detects endogenous levels of INT4 at Human/Mouse  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Integrator complex subunit 4 (Int4)  Observed Band 105kD  Cell Pathway Nucleus.  Tissue Specificity Aorta, Brain, Muscle, Ovary, Uterus,  Function function: Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (PCLR2A) and is recruited to the U1 and U2 snRNAs genes, similarity Belongs to the Integrator subunit 4 in INTS1, INTS3, INTS3, INTS3, INTS3, INTS3, INTS5, INTS6, INTS6	Catalog No	YP-mAb-08147
Applications WB Gene Name INTS4 MSTP093 Protein Name INT4 Immunogen Synthesized peptide derived from human INT4 AA range: 10-60 Specificity This antibody detects endogenous levels of INT4 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Integrator complex subunit 4 (Int4) Observed Band 105kD Cell Pathway Nucleus . Tissue Specificity Aorta, Brain, Muscle, Ovary, Uterus, Function function: Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 sRNAs genes, similarity belongs to the Integrator complex in Integrator to the Integrator usbunit 4 family, similarity Contains 8 HEAT repeats. subunit Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12., INTS6 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A) Mill 80660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; Mill 180680) do and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; Mill 180680) and U2 (RNU2; Mill 180680) (Baillat et al., 2005 [PubMed 162391444], [supplied by	Isotype	IgG
Gene Name         INTS4 MSTP093           Protein Name         INT4           Immunogen         Synthesized peptide derived from human INT4 AA range: 10-60           Specificity         This antibody detects endogenous levels of INT4 at Human/Mouse           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Integrator complex subunit 4 (Int4)           Observed Band         105kD           Cell Pathway         Nucleus .           Tissue Specificity         Aorta,Brain,Muscle,Ovary,Uterus,           Function         function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 sRNAs genes, similarity, Belongs to the Integrator to the Integrator complex in Integrator to the Integrator of the Integrator to the Integrator in Integrator in Integrator in Integrator in Integrator in Integrator in Integr	Reactivity	Human; Mouse
Immunogen   Synthesized peptide derived from human INT4 AA range: 10-60	Applications	WB
Immunogen         Synthesized peptide derived from human INT4 AA range: 10-60           Specificity         This antibody detects endogenous levels of INT4 at Human/Mouse           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Integrator complex subunit 4 (Int4)           Observed Band         105kD           Cell Pathway         Nucleus .           Tissue Specificity         Aorta, Brain, Muscle, Ovary, Uterus,           Function         function: Component of the Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes, similarity. Belongs to the Integrator subunit 4 family, similarity. Contains 8 HEAT repeats, subunit Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.           Background         Integrator complex subunit of the Integrator complex with the C-terminal domain (CRIVIZ; IMIM 180680) and III (RNU1; IMIM 180680) and III (RNU2; IMIM 180680) an	Gene Name	INTS4 MSTP093
Specificity         This antibody detects endogenous levels of INT4 at Human/Mouse           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Integrator complex subunit 4 (Int4)           Observed Band         105kD           Cell Pathway         Nucleus .           Tissue Specificity         Aorta,Brain,Muscle,Ovary,Uterus,           Function         function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes, similarity:Belongs to the Integrator subunit 4 family, similarity:Contains 8 HEAT repeats, subunit Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS5, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12, apprime and processing of small nuclear RNAs U1 (RNU1; MIM 180680) and u2 (RNU2; MIM 180680)	Protein Name	INT4
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Integrator complex subunit 4 (Int4)  Observed Band 105kD  Cell Pathway Nucleus .  Tissue Specificity Aorta,Brain,Muscle,Ovary,Uterus,  Function function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes, similarity:Belongs to the Integrator subunit 4 family, similarity:Contains 8 HEAT repeats. subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9, INTS	Immunogen	Synthesized peptide derived from human INT4 AA range: 10-60
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Integrator complex subunit 4 (Int4)         Observed Band       105kD         Cell Pathway       Nucleus .         Tissue Specificity       Aorta,Brain,Muscle,Ovary,Uterus,         Function       function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family,.similarity:Contains 8 HEAT repeats,.subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS9, INTS9	Specificity	This antibody detects endogenous levels of INT4 at Human/Mouse
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-1:2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Integrator complex subunit 4 (Int4)  Observed Band  105kD  Cell Pathway  Nucleus  Tissue Specificity  Aorta, Brain, Muscle, Ovary, Uterus,  function: Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes., similarity. Belongs to the Integrator subunit 4 family, similarity. Contains 8 HEAT repeats, subunit: Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background  INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillalt et al., 2005 [PubMed 16239144]). [supplied by	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.262% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Integrator complex subunit 4 (Int4)  Observed Band 105kD  Cell Pathway Nucleus .  Tissue Specificity Aorta,Brain,Muscle,Ovary,Uterus,  Function function:Component of the Integrator complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes, similarity:Belongs to the Integrator subunit 4 family, similarity:Contains 8 HEAT repeats, subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.  Background INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]),[supplied by	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Integrator complex subunit 4 (Int4)         Observed Band       105kD         Cell Pathway       Nucleus .         Tissue Specificity       Aorta,Brain,Muscle,Ovary,Uterus,         Function       function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of InTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12., INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12., Prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]), [supplied by	Purification	·
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms Integrator complex subunit 4 (Int4)  Observed Band 105kD  Cell Pathway Nucleus .  Tissue Specificity Aorta,Brain,Muscle,Ovary,Uterus,  Function function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A: MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Dilution	WB 1:500-1:2000
Synonyms Integrator complex subunit 4 (Int4)  Observed Band 105kD  Cell Pathway Nucleus .  Tissue Specificity Aorta,Brain,Muscle,Ovary,Uterus,  Function function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Concentration	1 mg/ml
Synonyms Integrator complex subunit 4 (Int4)  Observed Band 105kD  Cell Pathway Nucleus .  Tissue Specificity Aorta, Brain, Muscle, Ovary, Uterus,  Function function: Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes, similarity: Belongs to the Integrator subunit 4 family., similarity: Contains 8 HEAT repeats., subunit: Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]). [supplied by	Purity	≥90%
Observed Band  Cell Pathway  Nucleus .  Tissue Specificity  Aorta,Brain,Muscle,Ovary,Uterus,  function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background  INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Storage Stability	-20°C/1 year
Tissue Specificity  Aorta,Brain,Muscle,Ovary,Uterus,  function  function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background  INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Synonyms	Integrator complex subunit 4 (Int4)
Tissue Specificity  Aorta,Brain,Muscle,Ovary,Uterus,  function  function:Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background  INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Observed Band	105kD
function: Component of the Integrator complex, a complex involved in the small nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes., similarity: Belongs to the Integrator subunit 4 family., similarity: Contains 8 HEAT repeats., subunit: Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background  INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Cell Pathway	Nucleus .
nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes.,similarity:Belongs to the Integrator subunit 4 family.,similarity:Contains 8 HEAT repeats.,subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5, INTS6, INTS7, INTS8, INTS9/RC74, INTS10, CPSF3L/INTS11 and INTS12.,  Background  INTS4 is a subunit of the Integrator complex, which associates with the C-terminal domain of RNA polymerase II large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Tissue Specificity	Aorta,Brain,Muscle,Ovary,Uterus,
domain of RNA polymerase Il large subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by	Function	nuclear RNAs (snRNA) U1 and U2 transcription and in their 3'-box-dependent processing. The Integrator complex is associated with the C-terminal domain (CTD) of RNA polymerase II largest subunit (POLR2A) and is recruited to the U1 and U2 snRNAs genes., similarity:Belongs to the Integrator subunit 4 family., similarity:Contains 8 HEAT repeats., subunit:Belongs to the multiprotein complex Integrator, at least composed of INTS1, INTS2, INTS3, INTS4, INTS5,
	Background	domain of RNA polymerase Illarge subunit (POLR2A; MIM 180660) and mediates 3-prime end processing of small nuclear RNAs U1 (RNU1; MIM 180680) and U2 (RNU2; MIM 180690) (Baillat et al., 2005 [PubMed 16239144]).[supplied by



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using INT4 mouse mAb