




TBC16 mouse mAb

Catalog No	YP-mAb-08208
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TBC1D16
Protein Name	TBC16
Immunogen	Synthesized peptide derived from human TBC16 AA range: 529-579
Specificity	This antibody detects endogenous levels of TBC16 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.323% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1. mag/mal
Concentration	1 mg/ml
Purity	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Purity Storage Stability Synonyms	≥90% -20°C/1 year TBC1 domain family member 16
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year TBC1 domain family member 16 85kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year TBC1 domain family member 16 85kD early endosome,cytosol,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year TBC1 domain family member 16 85kD early endosome,cytosol, Brain,Breast,Skin,Urinary bladder, function:May act as a GTPase-activating protein for Rab family
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year TBC1 domain family member 16 85kD early endosome,cytosol, Brain,Breast,Skin,Urinary bladder, function:May act as a GTPase-activating protein for Rab family
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year TBC1 domain family member 16 85kD early endosome,cytosol, Brain,Breast,Skin,Urinary bladder, function:May act as a GTPase-activating protein for Rab family protein(s).,similarity:Contains 1 Rab-GAP TBC domain.,

Products Images



Western Blot analysis of various cells using TBC16 mouse mAb