





OR2H2 mouse mAb

| YP-mAb-08363 |
|--|
| IgG |
| Human;Rat;Mouse; |
| WB |
| OR2H2 FAT11 OLFR2 OR2H3 |
| OR2H2 |
| Synthesized peptide derived from human OR2H2 AA range: 240-290 |
| This antibody detects endogenous levels of OR2H2 at Human |
| Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Monoclonal, Mouse,IgG |
| The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| WB 1:500-1:2000 |
| 1 mg/ml |
| ≥90% |
| -20°C/1 year |
| |
| |
| Cell membrane; Multi-pass membrane protein. |
| |
| function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., |
| olfactory receptor family 2 subfamily H member 2(OR2H2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008], |
| |



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using OR2H2 mouse mAb