







KCC2G mouse mAb

Catalog No YP-mAb-08430 Isotype IgG Reactivity Human; Mouse;Rat	
Reactivity Human; Mouse;Rat	
Applications WB	
Gene Name CAMK2G CAMK CAMK-II CAMKG	
Protein Name KCC2G	
Immunogen Synthesized peptide derived from human KCC2G AA range: 490-540	
Specificity This antibody detects endogenous levels of KCC2G at Human/Mouse/Rat	t
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azid	le.
Source Monoclonal, Mouse,IgG	
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.	
Dilution WB 1:500-1:2000	
Concentration 1 mg/ml	
Purity ≥90%	
Storage Stability -20°C/1 year	
Synonyms	
Observed Band 61kD	
Cell Pathway Sarcoplasmic reticulum membrane ; Peripheral membrane protein ; Cytop side .	lasmic
Tissue Specificity Expressed in skeletal muscle.	
Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Autophosphorylation of CAMK2 plays an important role in the regulation of the kinase activity.,function:CaM-kinase II (CAMK2) is a pron kinase in the central nervous system that may function in long-term potent and neurotransmitter release.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. CAMK Suprotein kinase family. CaMK subfamily.,similarity:Contains 1 protein kinase domain.,subunit:CAMK2 is composed of four different chains: alpha, beta, gamma, and delta. The different isoforms assemble into homo- or heteromultimeric holoenzymes composed of 8 to 12 subunits.,	tiation er/Thr e
Background The product of this gene is one of the four subunits of an enzyme which to the serine/threonine protein kinase family, and to the Ca(2+)/calmodulin-dependent protein kinase subfamily. Calcium signaling crucial for several aspects of plasticity at glutamatergic synapses. In mam cells the enzyme is composed of four different chains: alpha, beta, gamma	is malian



UpingBio technology Co.,Ltd



delta. The product of this gene is a gamma chain. Many alternatively spliced transcripts encoding different isoforms have been described but the full-length nature of all the variants has not been determined.[provided by RefSeq, Mar


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using KCC2G mouse mAb