





ACOT6 mouse mAb

| Catalog No | YP-mAb-08551 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human; Mouse |
| Applications | WB |
| Gene Name | ACOT6 C14orf42 |
| Protein Name | ACOT6 |
| Immunogen | Synthesized peptide derived from human ACOT6 AA range: 116-166 |
| Specificity | This antibody detects endogenous levels of ACOT6 at Human/Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by |
| Dilution | affinity-chromatography using epitope-specific immunogen. WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | [Isoform 1]: Peroxisome . Localization to the peroxisome is uncertain since the potential C-terminal peroxisome targeting signal found in the mouse ortholog is not perfectly conserved; [Isoform 2]: Cytoplasm . Recombinant N-terminally GFP-tagged protein localizes to the cytosol |
| Tissue Specificity | |
| Function | caution:May lack catalytic activity, even though the active site residues are conserved.,function:Unknown.,similarity:Belongs to the C/M/P thioester hydrolase family., |
| Background | |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |














Western Blot analysis of various cells using ACOT6 mouse mAb