



VPRBP mouse mAb

Reactivity Human; Mouse Applications WB Gene Name VPRBP DCAF1 KIAA0800 RIP Protein Name VPRBP Immunogen Synthesized peptide derived from human VPRBP AA range: 804-854 Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis.		
Reactivity Human; Mouse Applications WB Gene Name VPRBP DCAF1 KIAA0800 RIP Protein Name VPRBP Immunogen Synthesized peptide derived from human VPRBP AA range: 804-854 Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain: Although not canonical, the WD repeat-like region is related to WD repeats, function: May act as a receptor for CUL4-DDB1 tubiquitin ligase complex. In case of infect to JU4-DDB2 using, it is ceruited by the VI por, possibly leading to the VPRBP family, similarity. Centains 1 Lisht domain, subunit: Interacts with PINY-1 virus Vpr protein and forms a tempary complex with HIV-1 Vpr and DDB1, tissue specificity. Ubiquitously expressed.,	Catalog No	YP-mAb-08619
Applications WB Gene Name VPRBP DCAF1 KIAA0800 RIP Protein Name VPRBP Immunogen Synthesized peptide derived from human VPRBP AA range: 804-854 Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner; recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats, function:May act as a receptor for CUL4-DDB1 ubiquitin liquise complex, in case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase, pathway: Protein modification; protein ubiquitination, similarity. Belongs to the VPRBP family, similarity. Contains 1 LisH domain. subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1, susse specificity. Ubiquitously expressed.,	Isotype	IgG
Gene Name VPRBP Protein Name VPRBP Immunogen Synthesized peptide derived from human VPRBP AA range: 804-854 Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner; recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. function domain:Although not canonical, the WD repeat-like region is related to WD repeats. function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 ry, possibly leading to hijack the RBX1-CUL4-DB2 and arrest the cell cycle in G2 phase, pathway:Protein modification, protein ubiquitinain, similarity:Belongs to the VPRBP family, similarity:Contains 1 LisH domain, subunit.Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1, tissue specificity:Ubiquitously expressed.	Reactivity	Human; Mouse
Protein Name VPRBP Immunogen Synthesized peptide derived from human VPRBP AA range: 804-854 Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats., function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus by r. possibly leading to hijack the RBX1-CUL4-DB2 and arrest the cell cycle in G2 phase, pathway. Protein modification; protein ubiquitination. similarity:Belongs to the VPRBP family. similarity:Contains 1 LisH domain., subunit.Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In	Applications	WB
Immunogen Synthesized peptide derived from human VPRBP AA range: 804-854 Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain: Although not canonical, the WD repeat-like region is related to WD repeats., function: May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase, pathway; Protein modification; protein ubiquitination, similarity:Belongs to the VPRBP family, similarity:Contains 1 LisH domain, subunit:Interacts with UPV-1 virus vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1., tissue specificity:Ubiquitously expressed. </td <td>Gene Name</td> <td>VPRBP DCAF1 KIAA0800 RIP</td>	Gene Name	VPRBP DCAF1 KIAA0800 RIP
Specificity This antibody detects endogenous levels of VPRBP at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before miltosis. Tissue Specificity Ubiquitously expressed. domain:Although not canonical, the WD repeat-like region is related to WD repeats, function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase, pathway.Protein modification, protein ubiquitination., similarity:Belongs to the VPRBP family, similarity:Contains 1 LisH domain., subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1, tissue specificity:Ubiquitously expressed.,	Protein Name	VPRBP
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cycle-dependent manner: recruited to chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before miltosis. Tissue Specificity Ubiquitously expressed. function domain:Although not canonical, the WD repeat-like region is related to WD repeats, function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase, pathway:Protein modification; protein ubiquitination, similarity:Belongs to the VPRBP family, similarity:Contains 1 Lish domain, subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.	Immunogen	Synthesized peptide derived from human VPRBP AA range: 804-854
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before miltosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats, function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase, pathway:Protein modification; protein ubiquitination, similarity. Belongs to the VPRBP family, similarity:Contains 1 LisH domain, subunit: Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1, tissue specificity:Ubiquitously expressed.,	Specificity	This antibody detects endogenous levels of VPRBP at Human/Mouse
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. domain:Although not canonical, the WD repeat-like region is related to WD repeats, function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase. pathway:Protein modification; protein ubiquitinationsimilarity:Belongs to the VPRBP family, similarity:Contains 1 LisH domain, subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a temary complex with HIV-1 Vpr and DDB1, tissue specificity:Ubiquitously expressed.,	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats, function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase, pathway:Protein modification; protein ubiquitination., similarity:Belongs to the VPRBP family, similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1, tissue specificity:Ubiquitously expressed.,	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Dilution	WB 1:500-1:2000
Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 Lish domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Purity	≥90%
Cell Pathway Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. Function domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase. pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Storage Stability	-20°C/1 year
Cytoplasm. Nucleus. Associated with chromatin in a DDB1-independent and cell cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Synonyms	
cycle-dependent manner: recruited to chromatin as DNA is being replicated and is released from chromatin before mitosis. Tissue Specificity Ubiquitously expressed. domain:Although not canonical, the WD repeat-like region is related to WD repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Observed Band	
domain:Although not canonical, the WD repeat-like region is related to WD repeats., function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase., pathway:Protein modification; protein ubiquitination., similarity:Belongs to the VPRBP family., similarity:Contains 1 LisH domain., subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1., tissue specificity:Ubiquitously expressed.,	Cell Pathway	cycle-dependent manner: recruited to chromatin as DNA is being replicated and is
repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1.,tissue specificity:Ubiquitously expressed.,	Tissue Specificity	Ubiquitously expressed.
Background	Function	repeats.,function:May act as a receptor for CUL4-DDB1 ubiquitin ligase complex. In case of infection by HIV-1 virus, it is recruited by HIV-1 Vpr, possibly leading to hijack the RBX1-CUL4-DDB2 and arrest the cell cycle in G2 phase.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the VPRBP family.,similarity:Contains 1 LisH domain.,subunit:Interacts with DDB1, suggesting that it may participate to some RBX1-CUL4-DDB2 ubiquitin ligase complex. In case of infection, interacts with HIV-1 virus Vpr protein and forms a ternary complex with HIV-1 Vpr and DDB1, tissue specificity:Ubiquitously
	Background	



UpingBio technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using VPRBP mouse mAb