






OSBP1 mouse mAb

Catalog No	YP-mAb-08716
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	OSBP OSBP1
Protein Name	OSBP1
Immunogen	Synthesized peptide derived from human OSBP1 AA range: 284-334
Specificity	This antibody detects endogenous levels of OSBP1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
- · ·	
Storage Stability	-20°C/1 year
	-20°C/1 year
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year Cytoplasm, cytosol . Cytoplasm, perinuclear region . Golgi apparatus membrane; Peripheral membrane protein . Endoplasmic reticulum membrane; Peripheral membrane protein . Golgi apparatus, trans-Golgi network . Predominantly cytosolic
Storage Stability Synonyms Observed Band	Cytoplasm, cytosol . Cytoplasm, perinuclear region . Golgi apparatus membrane; Peripheral membrane protein . Endoplasmic reticulum membrane; Peripheral membrane protein . Golgi apparatus, trans-Golgi network . Predominantly
Storage Stability Synonyms Observed Band Cell Pathway	Cytoplasm, cytosol . Cytoplasm, perinuclear region . Golgi apparatus membrane; Peripheral membrane protein . Endoplasmic reticulum membrane; Peripheral membrane protein . Golgi apparatus, trans-Golgi network . Predominantly cytosolic
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	Cytoplasm, cytosol . Cytoplasm, perinuclear region . Golgi apparatus membrane; Peripheral membrane protein . Endoplasmic reticulum membrane; Peripheral membrane protein . Golgi apparatus, trans-Golgi network . Predominantly cytosolic Widely expressed. domain:The PH and Ala/Gly-rich domains control cholesterol binding without affecting 25-hydroxycholesterol binding.,domain:The second coiled-coil domain is required for interaction with the tyrosine phosphatase.,function:Binds cholesterol and a range of oxysterols. Cholesterol binding promotes the formation of a complex with PP2A and a tyrosine phosphatase which dephosphorylate ERK1/2, whereas 25-hydroxycholesterol causes its disassembly. Regulates cholesterol efflux by decreasing ABCA1 stability.,similarity:Belongs to the OSBP family.,similarity:Contains 1 PH domain.,subcellular location:When bound to oxysterols, translocates to the periphery of Golgi membranes.,subunit:Homodimer




UpingBio technology Co.,Ltd








RefSeq, Jul 2008],



Products Images



Western Blot analysis of various cells using OSBP1 mouse mAb