



## SMOX mouse mAb

Catalog No	YP-mAb-08760
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	SMOX C20orf16 SMO UNQ3039/PRO9854
Protein Name	SMOX
Immunogen	Synthesized peptide derived from human SMOX AA range: 212-262
Specificity	This antibody detects endogenous levels of SMOX at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	[Isoform 1]: Cytoplasm. Nucleus.; [Isoform 4]: Cytoplasm. Nucleus.; [Isoform 6]: Cytoplasm. Nucleus.
Tissue Specificity	Widely expressed. Expressed in human tumor cell lines. Isoform 4 is only found in an embryonal kidney cell line.
Function	alternative products:At least 5 different isozymes are produced by alternative splicing of 8 exons. The resultant proteins have different biochemical characteristics and substrate specificity,catalytic activity:Spermidine + O(2) + H(2)O = putrescine + 3-aminopropanal + H(2)O(2).,catalytic activity:Spermine + O(2) + H(2)O = spermidine + 3-aminopropanal + H(2)O(2).,cofactor:Binds 1 FAD per subunit.,enzyme regulation:Inhibited at more than 90% by SL-11144, SL-11150 and SL-11158, at concentrations less than 1uM.,function:Flavoenzyme which catalyzes the oxidation of spermine to spermidine. Can also use N(1)-acetylspermine and spermidine as substrates, with different affinity depending on the isoform (isozyme) and on the experimental conditions. Plays an important role in the regulation of polyamine intracellular concentration and has the potential to act as a determinant of cellular sensitivi
Background	Polyamines are ubiquitous polycationic alkylamines which include spermine, spermidine, putrescine, and agmatine. These molecules participate in a broad range of cellular functions which include cell cycle modulation, scavenging



## UpingBio technology Co.,Ltd



reactive oxygen species, and the control of gene expression. These molecules also play important roles in neurotransmission through their regulation of cell-surface receptor activity, involvement in intracellular signalling pathways, and their putative roles as neurotransmitters. This gene encodes an FAD-containing enzyme that catalyzes the oxidation of spermine to spermadine and secondarily produces hydrogen peroxide. Multiple transcript variants encoding different isoenzymes have been identified for this gene, some of which have failed to demonstrate significant oxidase activity on natural polyamine substrates. The characterized isoenzymes have distinctive biochemical characteristics and substrate specificities, suggesting the existence of additional levels of complexity in polyamine catabolism. [provided by RefSeq, Jul 2012],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using SMOX mouse mAb