



NLTP mouse mAb

Catalog No	YP-mAb-08965
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	SCP2
Protein Name	NLTP
Immunogen	Synthesized peptide derived from human NLTP AA range: 338-388
Specificity	This antibody detects endogenous levels of NLTP at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	[Isoform SCP2]: Peroxisome . Cytoplasm . Mitochondrion . Endoplasmic reticulum . Mitochondrion .; [Isoform SCPx]: Peroxisome .
Tissue Specificity	Liver, fibroblasts, and placenta.
Function	catalytic activity:3-alpha,7-alpha,12-alpha-trihydroxy-5-beta-cholanoyl-CoA + propanoyl-CoA = CoA + 3-alpha,7-alpha,12-alpha-trihydroxy-24-oxo-5-beta-cholestanoyl-CoA.;caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.;disease:SCP2 is present in low levels in subjects with Zellweger syndrome (cerebro-hepatic-renal syndrome), whose cells are deficient in peroxisomes and who have an associated impairment in plasmalogen and bile acid synthesis and catabolism of phytanic acid and very-long-chain fatty acids.;function:Mediates in vitro the transfer of all common phospholipids, cholesterol and gangliosides between membranes. May play a role in regulating steroidogenesis.;similarity:Belongs to the thiolase family.;similarity:Contains 1 SCP2 domain.;similarity:In the N-terminal section; belongs to the thiolase fami
Background	This gene encodes two proteins: sterol carrier protein X (SCPx) and sterol carrier protein 2 (SCP2), as a result of transcription initiation from 2 independently



regulated promoters. The transcript initiated from the proximal promoter encodes the longer SCPx protein, and the transcript initiated from the distal promoter encodes the shorter SCP2 protein, with the 2 proteins sharing a common C-terminus. Evidence suggests that the SCPx protein is a peroxisome-associated thiolase that is involved in the oxidation of branched chain fatty acids, while the SCP2 protein is thought to be an intracellular lipid transfer protein. This gene is highly expressed in organs involved in lipid metabolism, and may play a role in Zellweger syndrome, in which cells are deficient in peroxisomes and have impaired bile acid synthesis. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms.[provided by RefSeq, Aug 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using NLTP mouse mAb