





O5AL1 mouse mAb

Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity function:Odorant receptor "polymorphism:A double nucleotide deletion at procession of the protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene., similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes are the genes are fully to the olfactory receptor genes are the genes are fully to the olfactory receptor genes are the genes are fully to the olfactory receptor genes are the genes are fully to the olfactory receptor genes are the genes are fully to the olfactory receptor genes are fully to the processor of th		
Reactivity Human;Rat;Mouse; Applications WB Gene Name OR5AL1 OR5AL1P Protein Name O5AL1 Immunogen Synthesized peptide derived from human O5AL1 AA range: 131-181 Specificity This antibody detects endogenous levels of O5AL1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor ,polymorphism:A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene, similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptors (GP arising from single coding-exon genes. Olfactory receptors family is the lar the genome. The nomenclature assigned to the olfactory receptor gene family is the lar the genome. The nomenclature assigned to the lolfactory receptor gene family is the lar the genome. The nomenclature assigned to the lolfactory receptor gene family is the lar the genome. The nomenclature assigned to the lolfactory receptor gene family is the lar the genome. The nomenclature assigned to the lolfactory receptor gene family of the protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the lolfactory receptor genes and are responsible for the receptor genes and are responsible for the receptor genes and are responsible for	Catalog No	YP-mAb-09122
Applications WB Gene Name OR5AL1 OR5AL1P Protein Name O5AL1 Immunogen Synthesized peptide derived from human O5AL1 AA range: 131-181 Specificity This antibody detects endogenous levels of O5AL1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor polymorphism: A double nucleotide deletion at procys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene, similarity. Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors garder in the genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes and are responsible for the recognition and G protein-mediated	Isotype	IgG
Gene Name OR5AL1 OR5AL1P Protein Name O5AL1 Immunogen Synthesized peptide derived from human O5AL1 AA range: 131-181 Specificity This antibody detects endogenous levels of O5AL1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptorpolymorphism: A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene, similarity: Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are an 7-transmembrane domain structure with many neuronasmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome. The nomenclature assigned to the olfactory receptor genes and the qenome	Reactivity	Human;Rat;Mouse;
Protein Name O5AL1 Immunogen Synthesized peptide derived from human O5AL1 AA range: 131-181 Specificity This antibody detects endogenous levels of O5AL1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptorpolymorphism: A double nucleotide deletion at presponsible for functional diversity thus producing a pseudogene, similarity: Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors GP- arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes and the genome.	Applications	WB
Immunogen Synthesized peptide derived from human O5AL1 AA range: 131-181 Specificity This antibody detects endogenous levels of O5AL1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity function: Odorant receptorpolymorphism: A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene;similarity:Belongs to the G-protein coupled receptor 1 family pseudogene;similarity:Belongs to the G-protein coupled receptor 1 family protein responsible for functional diversity thus producing a pseudogene;similarity:Belongs to the G-protein coupled receptors of a protein receptor gene and a protein-coupled receptors of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. T	Gene Name	OR5AL1 OR5AL1P
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptorpolymorphism: A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogenesimilarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes and the genome. The nomenclature assigned to the olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes are the genome. The nomenclature assigned to the olfactory receptor genes are	Protein Name	O5AL1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptorpolymorphism: A double nucleotide deletion at process companies of the protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene, similarity: Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GP) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes are the genome. The nomenclature assignade to the olfactory receptor genes are the genome. The nomenclature assignated to the olfactory receptor genes are the genome. The nomenclature assignade to the olfactory receptor genes are	Immunogen	Synthesized peptide derived from human O5AL1 AA range: 131-181
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptorpolymorphism:A double nucleotide deletion at processors of the protein produces a stop codon which is responsible for functional diversity thus producing a pseudogenesimilarity:Belongs to the G-protein coupled receptor 1 family Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the quenome. The nomencladure assigned to the olfactory receptor genes are the quenome. The nomencladure assigned to the lofactory receptor genes are the quenome. The nomencladure assigned to the lofactory receptor genes are the quenome. The nomencladure assigned to the lofactory receptor genes are the percention of the protein coupled receptor genes are the protein coupled receptor genes are the protein coupled receptor genes are the protein coup	Specificity	This antibody detects endogenous levels of O5AL1 at Human
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor ,,polymorphism:A double nucleotide deletion at procys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptores are members of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptors (GP arising from single coding-exon genes. Olfactory receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-1:2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,polymorphism:A double nucleotide deletion at process a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptoris are members of a large family of G-protein-coupled receptors (GP) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes family is the lar the genome. The nomenclature assigned to the olfactory receptor genes are	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptorpolymorphism:A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes are	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,polymorphism:A double nucleotide deletion at process a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptoreins are members of a large family of G-protein-coupled receptors (GP-arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lart the genome. The nomenclature assigned to the olfactory receptor genes are	Dilution	WB 1:500-1:2000
Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor .,polymorphism: A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GParising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes are	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,polymorphism:A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GParising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormore receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lart the genome. The nomenclature assigned to the olfactory receptor genes are	Purity	≥90%
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,polymorphism:A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GParising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lart the genome. The nomenclature assigned to the olfactory receptor genes are	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function: Odorant receptor .,polymorphism: A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lart the genome. The nomenclature assigned to the olfactory receptor genes are	Synonyms	
Function function:Odorant receptor .,polymorphism:A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptorins are members of a large family of G-protein-coupled receptors (GPG arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormoreceptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lart the genome. The nomenclature assigned to the olfactory receptor genes are	Observed Band	
Function function:Odorant receptor .,polymorphism:A double nucleotide deletion at p Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory rec proteins are members of a large family of G-protein-coupled receptors (GP- arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes ar	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene., similarity: Belongs to the G-protein coupled receptor 1 family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GP arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes are	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory rec proteins are members of a large family of G-protein-coupled receptors (GP) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormo receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the lar the genome. The nomenclature assigned to the olfactory receptor genes ar	Function	function:Odorant receptor .,polymorphism:A double nucleotide deletion at position Cys-156 in the gene coding for this protein produces a stop codon which is responsible for functional diversity thus producing a pseudogene.,similarity:Belongs to the G-protein coupled receptor 1 family.,
RefSeq, Jul 2008],	Background	7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using O5AL1 mouse mAb