





PRAF3 mouse mAb


| Catalog No | YP-mAb-09124 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human; Mouse;Rat |
| Applications | WB |
| Gene Name | ARL6IP5 DERP11 JWA PRA2 PRAF3 HSPC127 |
| Protein Name | PRAF3 |
| Immunogen | Synthesized peptide derived from human PRAF3 AA range: 120-170 |
| Specificity | This antibody detects endogenous levels of PRAF3 at Human/Mouse/Rat |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-1:2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | Endoplasmic reticulum membrane ; Multi-pass membrane protein . Cell membrane ; Multi-pass membrane protein . Cytoplasm . Cytoplasm, cytoskeleton . Also exists as a soluble form in the cytoplasm. Associated with microtubules |
| Tissue Specificity | |
| Function | function:Regulates intracellular concentrations of taurine and glutamate. |

Background

Expression of this gene is affected by vitamin A. The encoded protein of this gene may be associated with the cytoskeleton. A similar protein in rats may play a role in the regulation of cell differentiation. The rat protein binds and inhibits the cell membrane glutamate transporter EAAC1. The expression of the rat gene is



UpingBio technology Co.,Ltd







upregulated by retinoic acid, which results in a specific reduction in EAAC1-mediated glutamate transport. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PRAF3 mouse mAb