


HACD3 mouse mAb

Catalog No	YP-mAb-09156
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	PTPLAD1 BIND1 HACD3
Protein Name	HACD3
Immunogen	Synthesized peptide derived from human HACD3 AA range: 255-305
Specificity	This antibody detects endogenous levels of HACD3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Band Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Highly expressed in testis, kidney, brain, liver and weakly in skeletal muscle, spleen and heart. No expression detected in leukocytes.
Cell Pathway	Highly expressed in testis, kidney, brain, liver and weakly in skeletal muscle, spleen and heart. No expression detected in leukocytes. function:Involved in Rac1-signaling pathways leading to the modulation of gene expression, induction:By AKAP12 and histone deacetylase inhibitors such as
Cell Pathway Tissue Specificity	Highly expressed in testis, kidney, brain, liver and weakly in skeletal muscle, spleen and heart. No expression detected in leukocytes. function:Involved in Rac1-signaling pathways leading to the modulation of gene expression.,induction:By AKAP12 and histone deacetylase inhibitors such as sodium butyrate.,similarity:Belongs to the PTPLA family.,similarity:Contains 1 CS domain.,subunit:Interacts with RAC1.,tissue specificity:Highly expressed in testis,
Cell Pathway Tissue Specificity Function	Highly expressed in testis, kidney, brain, liver and weakly in skeletal muscle, spleen and heart. No expression detected in leukocytes. function:Involved in Rac1-signaling pathways leading to the modulation of gene expression.,induction:By AKAP12 and histone deacetylase inhibitors such as sodium butyrate.,similarity:Belongs to the PTPLA family.,similarity:Contains 1 CS domain.,subunit:Interacts with RAC1.,tissue specificity:Highly expressed in testis,
Cell Pathway Tissue Specificity Function Background matters needing	Highly expressed in testis, kidney, brain, liver and weakly in skeletal muscle, spleen and heart. No expression detected in leukocytes. function:Involved in Rac1-signaling pathways leading to the modulation of gene expression.,induction:By AKAP12 and histone deacetylase inhibitors such as sodium butyrate.,similarity:Belongs to the PTPLA family.,similarity:Contains 1 CS domain.,subunit:Interacts with RAC1.,tissue specificity:Highly expressed in testis, kidney, brain, liver and weakly in skeletal muscle, spleen and heart.,



Products Images



Western Blot analysis of various cells using HACD3 mouse mAb