





## NLRX1 Monoclonal Antibody

| Catalog No         | YP-mAb-10611   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Mouse;Rat  |
| Applications       | WB   |
| Gene Name          | NLRX1  |
| Protein Name       | NLRX1  |
| lmmunogen          | The antiserum was produced against synthesized peptide derived from the Internal region of human NLRX1. AA range:581-630   |
| Specificity        | NLRX1 Monoclonal Antibody detects endogenous levels of NLRX1   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Monoclonal, Mouse,IgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-1:2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | NLR family member X1 (Caterpiller protein 11.3;CLR11.3;Nucleotide-binding oligomerization domain protein 26;Nucleotide-binding oligomerization domain protein 5;Nucleotide-binding oligomerization domain protein 9)   |
| Observed Band      | 108kD  |
| Cell Pathway       | Mitochondrion outer membrane .   |
| Tissue Specificity | Ubiquitously expressed. Strongest expression in mammary gland, heart and muscle. Detected in HeLa, HEK293T, THP-1, HL-60, Raji and Jurkat cell lines (at protein level).   |
| Function           | function:Participates in antiviral signaling. Acts as a negative regulator of MAVS-mediated antiviral responses, through the inhibition of the virus-induced RLH (RIG-like helicase)-MAVS interaction (PubMed:18200010). Has no inhibitory function on NF-Kappa-B and type 1 interferon signaling pathways, but enhances NF-Kappa-B and JUN N-terminal kinase dependent signaling through the production of reactive oxygen species (PubMed:18219313).,sequence caution:Translated as Trp.,sequence caution:Translation N-terminally extended.,similarity:Belongs to the NLRP family.,similarity:Contains 1 NACHT domain.,similarity:Contains 4 LRR (leucine-rich) repeats.,subunit:Interacts with MAVS.,tissue specificity:Ubiquitously expressed. Strongest expression in mammary gland, heart and muscle. Detected in HELA, 293T, THP-1, HL60, RAJI, and JURKAT cell lines (at protein level)., |



## UpingBio technology Co.,Ltd






| Background      | The protein encoded by this gene is a member of the NLR family and localizes to the outer mitochondrial membrane. The encoded protein is a regulator of mitochondrial antivirus responses. Three transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Aug 2013], |
|-----------------|---|
| matters needing | Avoid repeated freezing and thawing!  |

attention

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. **Usage suggestions** 

## **Products Images**



Western Blot analysis of various cells using NLRX1 Monoclonal Antibody