



GP-39 Monoclonal Antibody

Catalog No	YP-mAb-10860
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CHI3L1
Protein Name	GP-39
Immunogen	Synthesized peptide derived from human GP-39
Specificity	This antibody detects endogenous levels of human GP-39
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Chitinase-3-like protein 1 (39 kDa synovial protein;Cartilage glycoprotein 39;CGP-39;GP-39;hCGP-39;YKL-40)
Observed Band	42kD
Cell Pathway	Secreted, extracellular space . Cytoplasm . Cytoplasm, perinuclear region . Endoplasmic reticulum .
Tissue Specificity	Present in activated macrophages, articular chondrocytes, synovial cells as well as in liver. Very low or undetectable expression in non-inflammatory colon. Undetectable in muscle tissues, lung, pancreas, mononuclear cells, or fibroblasts.
Function	disease:A genetic variation in CHI3L1 is associated with susceptibility to asthma-related traits type 7 (ASRT7) [MIM:611960]. Asthma-related traits include clinical symptoms of asthma, such as coughing, wheezing and dyspnea, bronchial hyperresponsiveness (BHR) as assessed by methacholine challenge test, serum IgE levels, atopy, and atopic dermatitis.,function:Carbohydrate-binding lectin with a preference for chitin. May play a role in defense against pathogens, or in tissue remodeling. May play an important role in the capacity of cells to respond to and cope with changes in their environment.,PTM:Glycosylated.,similarity:Belongs to the glycosyl hydrolase 18 family.,subunit:Monomer.,tissue specificity:Present in activated macrophages, articular chondrocytes, synovial cells as well as in liver. Undetectable in muscle tissues, lung, pancreas, mononuclear cells, or fibroblasts.,



Background

Chitinases catalyze the hydrolysis of chitin, which is an abundant glycopolymer found in insect exoskeletons and fungal cell walls. The glycoside hydrolase 18 family of chitinases includes eight human family members. This gene encodes a glycoprotein member of the glycosyl hydrolase 18 family. The protein lacks chitinase activity and is secreted by activated macrophages, chondrocytes, neutrophils and synovial cells. The protein is thought to play a role in the process of inflammation and tissue remodeling. [provided by RefSeq, Sep 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GP-39 Monoclonal Antibody