



T2H2L mouse mAb

YP-mAb-10976
IgG
Human;Rat;Mouse;
WB
GTF2H2C; GTF2H2D
T2H2L
Synthesized peptide derived from human T2H2L AA range: 275-325
This antibody detects endogenous levels of T2H2L at Human
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
Nucleus .
alternative products: A number of isoforms may be produced. The isoforms may be also produced by incomplete gene duplication, function: Component of the core-TFIIH basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II., function: Component of the core-TFIIH basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. The N-terminus interacts with and regulates XPD whereas an intact C-terminus is required for a successful escape of RNAP II form the promoter., similarity: Belongs to the GTF2H2 family., similarity: Contains 1 VWFA domain., subunit: One of the six subunits forming the core-TFIIH basal transcription factor. Interacts with XPB, XPD, GTF2H1 and GTF2H3., tissue specificity: Widely expressed, with higher express

Background



UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using T2H2L mouse mAb