





LRWD1 mouse mAb

Catalog No	YP-mAb-11017
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	LRWD1 ORCA
Protein Name	LRWD1
Immunogen	Synthesized peptide derived from human LRWD1 AA range: 395-445
Specificity	This antibody detects endogenous levels of LRWD1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus . Chromosome, centromere. Chromosome, telomere. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Chromosome, centromere, kinetochore . Localizes to heterochromatin during G1 phase. Restricted to centromeres or telomeres as cells progress though S phase. When cells enter mitosis, relocalizes to centromeres. Recruitment to pericentric heterochromatin largely depends on the presence of H3K9me3.
Cell Pathway Tissue Specificity	cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Localizes to heterochromatin during G1 phase. Restricted to centromeres or telomeres as cells progress though S phase. When cells enter mitosis, relocalizes to centromeres. Recruitment to pericentric
-	cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Localizes to heterochromatin during G1 phase. Restricted to centromeres or telomeres as cells progress though S phase. When cells enter mitosis, relocalizes to centromeres. Recruitment to pericentric heterochromatin largely depends on the presence of H3K9me3. Testis-specific. Drastically down-regulated in testis from patients with Sertoli
Tissue Specificity	cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Localizes to heterochromatin during G1 phase. Restricted to centromeres or telomeres as cells progress though S phase. When cells enter mitosis, relocalizes to centromeres. Recruitment to pericentric heterochromatin largely depends on the presence of H3K9me3. Testis-specific. Drastically down-regulated in testis from patients with Sertoli cell-only syndrome (SCOS). PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 3 LRR (leucine-rich) repeats.,similarity:Contains 5 WD



UpingBio technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using LRWD1 mouse mAb