



ICT1 mouse mAb

| YP-mAb-11056 |
|---|
| IgG |
| Human; Mouse |
| WB |
| ICT1 DS1 |
| ICT1 |
| Synthesized peptide derived from human ICT1 AA range: 7-57 |
| This antibody detects endogenous levels of ICT1 at Human/Mouse |
| Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Monoclonal, Mouse,IgG |
| The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| WB 1:500-1:2000 |
| 1 mg/ml |
| ≥90% |
| -20°C/1 year |
| |
| |
| Mitochondrion . |
| Down-regulated during the in vitro differentiation of HT29-D4 colon carcinoma cells. |
| similarity:Belongs to the prokaryotic/mitochondrial release factor family.,tissue specificity:Down-regulated during the in vitro differentiation of HT29-D4 colon carcinoma cells., |
| The protein encoded by this gene is a peptidyl-tRNA hydrolase and a vital component of the large mitochondrial ribosome. The encoded protein serves as a ribosome release factor for this ribosome, which translates mitochondrial genes. This protein may be responsible for degrading prematurely-terminated polypeptides and for reusing stalled ribosomes. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2014], |
| Avoid repeated freezing and thawing! |
| |



UpingBio technology Co.,Ltd







Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



