





## SPRE2 mouse mAb

Reactivity       Human; Mouse;Rat         Applications       WB         Gene Name       SPRED2         Protein Name       SPRE2         Immunogen       Synthesized peptide derived from human SPRE2 AA range: 68-118         Specificity       This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse, lgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cell membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, Secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Catalog No	YP-mAb-11451
Applications  WB  Gene Name SPRED2  Protein Name SPRE2  Immunogen Synthesized peptide derived from human SPRE2 AA range: 68-118  Specificity This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side . Cytoplasmic side . Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Isotype	IgG
Gene Name         SPRED2           Protein Name         SPRE2           Immunogen         Synthesized peptide derived from human SPRE2 AA range: 68-118           Specificity         This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms           Observed Band         Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic side. Cytoplasmic vesicles that are supposed to be secretory granules.	Reactivity	Human; Mouse;Rat
Protein Name         SPRE2           Immunogen         Synthesized peptide derived from human SPRE2 AA range: 68-118           Specificity         This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-1:2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms           Observed Band         Cell Pathway           Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Stroplasmic side . Cytoplasmic si	Applications	WB
Immunogen       Synthesized peptide derived from human SPRE2 AA range: 68-118         Specificity       This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cell membrane; Peripheral membrane protein; Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic vesicle Cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules	Gene Name	SPRED2
Specificity       This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Monoclonal, Mouse, IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cell membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Protein Name	SPRE2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle secretory vesicle membrane ; Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Immunogen	Synthesized peptide derived from human SPRE2 AA range: 68-118
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-1:2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band         Cell Pathway       Cell membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, side . Cytoplasmic vesicles that are supposed to be secretory granules	Specificity	This antibody detects endogenous levels of SPRE2 at Human/Mouse/Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic side . Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic side . Cytoplasmic vesicle stat are supposed to be secretory granules	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-1:2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic vesicles that are supposed to be secretory granules	Source	Monoclonal, Mouse,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cell membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic vesicle, secretory vesicle in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle. Cytoplasm. Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Dilution	WB 1:500-1:2000
Storage Stability  Synonyms  Observed Band  Cell Pathway  Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle . Cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules	Concentration	1 mg/ml
Synonyms  Observed Band  Cell Pathway  Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm . Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules	Purity	≥90%
Observed Band  Cell Pathway  Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm . Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasmic vesicle, secretory vesicle membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm . Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules	Synonyms	
vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm. Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory granules.	Observed Band	
<b>Tissue Specificity</b> Expressed in liver, skin, small intestine, salivary gland and prostate.	Cell Pathway	vesicle, secretory vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm. Detected in the cytoplasm of the stratum spinosum cells, where it is associated with cytoplasmic vesicles that are supposed to be secretory
	Tissue Specificity	Expressed in liver, skin, small intestine, salivary gland and prostate.
	Function	of MAP kinase.,PTM:Phosphorylated on tyrosine. Phosphorylation of Tyr-228 and Tyr-231 are required for ubiquitination.,PTM:Ubiquitinated; leading to degradation by the proteasome.,similarity:Contains 1 KBD domain.,similarity:Contains 1 SPR (sprouty) domain.,similarity:Contains 1 WH1 domain.,subcellular location:Especially found in the cytoplasm of the stratum spinosum cells called membrane-coated granules.,subunit:Interacts with Ras (By similarity). Homodimer and heterodimer. Able to interact with SPRED1 to form heterodimers.,tissue specificity:Expressed in liver, skin, small intestine, salivary
Background SPRED2 is a member of the Sprouty (see SPRY1; MIM 602465)/SPRED family of proteins that regulate growth factor-induced activation of the MAP kinase cascades.	Background	SPRED2 is a member of the Sprouty (see SPRY1; MIM 602465)/SPRED family of proteins that regulate growth factor-induced activation of the MAP kinase cascade



## UpingBio technology Co.,Ltd



(see MAPK1; MIM 176948) (Nonami et al., 2004 [PubMed 15465815]).[supplied by OMIM, Mar 2008],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using SPRE2 mouse mAb