







ZEP2 mouse mAb

YP-mAb-11497
IgG
Human; Mouse;Rat
WB
HIVEP2
ZEP2
Synthesized peptide derived from human ZEP2 AA range: 1975-2025
This antibody detects endogenous levels of ZEP2 at Human/Mouse/Rat
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
Nucleus.
Expressed in brain and skeletal muscle.
function:This protein specifically binds to the DNA sequence 5'-GGGACTTTCC-3' which is found in the enhancer elements of numerous viral promoters such as those of SV40, CMV, or HIV1. In addition, related sequences are found in the enhancer elements of a number of cellular promoters, including those of the class I MHC, interleukin-2 receptor, somatostatin receptor II, and interferon-beta genes. It may act in T-cell activation.,induction:By mitogens and phorbol ester.,similarity:Contains 4 C2H2-type zinc fingers.,subunit:Interacts with TCF4.,tissue specificity:Expressed in brain and skeletal muscle.,
This gene encodes a member of a family of closely related, large, zinc finger-containing transcription factors. The encoded protein regulates transcription by binding to regulatory regions of various cellular and viral genes that maybe involved in growth, development and metastasis. The protein contains the ZAS domain comprised of two widely separated regions of zinc finger motifs, a stretch of highly acidic amino acids and a serine/threonine-rich sequence. [provided by



UpingBio technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using ZEP2 mouse mAb