





## LX15B mouse mAb

Catalog No	YP-mAb-11594
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	ALOX15B
Protein Name	LX15B
Immunogen	Synthesized peptide derived from human LX15B AA range: 576-626
Specificity	This antibody detects endogenous levels of LX15B at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	74kD
Cell Pathway	[Isoform A]: Nucleus . Other isoforms are excluded from the nucleus; Cytoplasm, cytosol . Cell membrane . Cytoplasm, cytoskeleton . Membrane ; Peripheral membrane protein . Cell junction, adherens junction . Cell junction, focal adhesion . Predominantly cytosolic; becomes enriched at membranes upon calcium binding
Tissue Specificity	Expressed in hair, prostate, lung, ovary, lymph node, spinal cord and cornea.
Function	catalytic activity:Arachidonate + O(2) = (5Z,8Z,11Z,13E)-(15S)-15-hydroperoxyicosa-5,8,11,13-tetraenoate.,cofactor:Binds 1 iron ion per subunit.,function:Converts arachidonic acid exclusively to 15S-hydroperoxyeicosatetraenoic acid, while linoleic acid is less well metabolized.,pathway:Lipid metabolism; leukotriene D4 biosynthesis.,similarity:Belongs to the lipoxygenase family.,similarity:Contains 1 lipoxygenase domain.,similarity:Contains 1 PLAT domain.,tissue specificity:Expressed in hair, prostate, lung and cornea.,
Background	This gene encodes a member of the lipoxygenase family of structurally related nonheme iron dioxygenases involved in the production of fatty acid hydroperoxides. The encoded protein converts arachidonic acid exclusively to 15S-hydroperoxyeicosatetraenoic acid, while metabolizing linoleic acid less



## UpingBio technology Co.,Ltd





effectively. This gene is located in a cluster of related genes and a pseudogene that spans approximately 100 kilobases on the short arm of chromosome 17. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using LX15B mouse mAb