







## RPIA mouse mAb

| Catalog No         | YP-mAb-11617   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human; Mouse   |
| Applications       | WB   |
| Gene Name          | RPIA RPI   |
| Protein Name       | RPIA   |
| Immunogen          | Synthesized peptide derived from human RPIA AA range: 256-306  |
| Specificity        | This antibody detects endogenous levels of RPIA at Human/Mouse   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Monoclonal, Mouse,IgG  |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-1:2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           |  |
| Observed Band      |  |
| Cell Pathway       | cytosol,integral component of membrane,intracellular membrane-bounded organelle,   |
| Tissue Specificity |  |
| Function           | catalytic activity:D-ribose 5-phosphate = D-ribulose 5-phosphate.,disease:Defects in RPIA are the cause of ribose 5-phosphate isomerase deficiency [MIM:608611]. A patient has been described with a deficiency of ribose 5-phosphate isomerase who presented with leukoencephalopathy and peripheral neuropathy. Proton magnetic resonance spectroscopy of the brain revealed a highly elevated level of the polyols ribitol and D-arabitol, which were subsequently also found in high concentrations in body fluids. Deficient activity of RPIA, one of the pentose phosphate pathway enzymes, has been demonstrated in fibroblasts.,pathway:Carbohydrate degradation; pentose phosphate pathway; D-ribose 5-phosphate from D-ribulose 5-phosphate (non-oxidative stage): step 1/1.,similarity:Belongs to the ribose 5-phosphate isomerase family., |
| Background         | The protein encoded by this gene is an enzyme, which catalyzes the reversible conversion between ribose-5-phosphate and ribulose-5-phosphate in the pentose-phosphate pathway. This gene is highly conserved in most organisms. The enzyme plays an essential role in the carbohydrate metabolism. Mutations in  |



## UpingBio technology Co.,Ltd







this gene cause ribose 5-phosphate isomerase deficiency. A pseudogene is found on chromosome 18. [provided by RefSeq, Mar 2010],


| matters needing |  |
|-----------------|--|
| attention       |  |

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using RPIA mouse mAb