



PRDM2 mouse mAb

Catalog No	YP-mAb-11653
Isotype	IgG
Reactivity	Human;Rat
Applications	WB
Gene Name	PRDM2 KMT8 RIZ
Protein Name	PRDM2
Immunogen	Synthesized peptide derived from human PRDM2 AA range: 964-1014
Specificity	This antibody detects endogenous levels of PRDM2 at Human/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus .
Tissue Specificity	Highly expressed in retinoblastoma cell lines and in brain tumors. Also expressed in a number of other cell lines and in brain, heart, skeletal muscle, liver and spleen. Isoform 1 is expressed in testis at much higher level than isoform 3.
Function	catalytic activity:S-adenosyl-L-methionine + histone L-lysine = S-adenosyl-L-homocysteine + histone N(6)-methyl-L-lysine.;function:S-adenosyl-L-methionine-dependent histone methyltransferase that specifically methylates 'Lys-9' of histone H3. May function as a DNA-binding transcription factor. Binds to the macrophage-specific TPA-responsive element (MTE) of the HMOX1 (heme oxygenase 1) gene and may act as a transcriptional activator of this gene.;similarity:Contains 1 SET domain.;similarity:Contains 8 C2H2-type zinc fingers.;subunit:Binds to the retinoblastoma protein (RB). Interacts with GATA3.;tissue specificity:Highly expressed in retinoblastoma cell lines and in brain tumors. Also expressed in a number of other cell lines and in brain, heart, skeletal muscle, liver and spleen. Isoform 1 is expressed in testis at much higher level than isoform 3.,
Background	This tumor suppressor gene is a member of a nuclear histone/protein methyltransferase superfamily. It encodes a zinc finger protein that can bind to



retinoblastoma protein, estrogen receptor, and the TPA-responsive element (MTE) of the heme-oxygenase-1 gene. Although the functions of this protein have not been fully characterized, it may (1) play a role in transcriptional regulation during neuronal differentiation and pathogenesis of retinoblastoma, (2) act as a transcriptional activator of the heme-oxygenase-1 gene, and (3) be a specific effector of estrogen action. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PRDM2 mouse mAb