





## EXOS1 mouse mAb

| Catalog No         | YP-mAb-11666  |
|--------------------|---|
| Isotype            | IgG   |
| Reactivity         | Human; Mouse  |
| Applications       | WB  |
| Gene Name          | EXOSC1 CSL4 CGI-108   |
| Protein Name       | EXOS1   |
| Immunogen          | Synthesized peptide derived from human EXOS1 AA range: 93-143   |
| Specificity        | This antibody detects endogenous levels of EXOS1 at Human/Mouse   |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source             | Monoclonal, Mouse,IgG   |
| Purification       | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution           | WB 1:500-1:2000   |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           |   |
| Observed Band      |   |
| Cell Pathway       | Nucleus, nucleolus . Nucleus . Cytoplasm .  |
| Tissue Specificity |   |
| Function           | domain:The N- and C-terminal regions of the protein are required for interaction with RRP46.,function:Component of the exosome 3'->5' exoribonuclease complex, a complex that degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions. Required for the 3'processing of the 7S pre-RNA to the mature 5.8S rRNA. Has a 3'-5' exonuclease activity.,similarity:Contains 1 S1 motif domain.,subunit:Component of the exosome multienzyme ribonuclease complex composed of at least 11 proteins: EXOSC1/CSL4, EXOSC2/RRP4, EXOSC3/RRP40, EXOSC4/RRP41, EXOSC5/RRP46, EXOSC6/MTR3, EXOSC7/RRP42, EXOSC8/RRP43, EXOSC9/ScI-75, DIS3/RRP44, and EXOSC10/ScI-100 (only in the nuclear complex). Interacts directly with EXOSC5, EXOSC6, EXOSC7 and EXOSC8. Also associated with the GTPase Ran., |
| Background         | This gene encodes a core component of the exosome. The mammalian exosome is required for rapid degradation of AU rich element-containing RNAs but not for poly(A) shortening. The association of this protein with the exosome is mediated  |



## UpingBio technology Co.,Ltd



by protein-protein interactions with ribosomal RNA-processing protein 42 and ribosomal RNA-processing protein 46. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Jan 2016],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using EXOS1 mouse mAb