




## TM192 mouse mAb

Catalog No	YP-mAb-11844
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TMEM192
Protein Name	TM192
Immunogen	Synthesized peptide derived from human TM192 AA range: 27-77
Specificity	This antibody detects endogenous levels of TM192 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	4 manufactural
Concentration	1 mg/ml
Purity	1 mg/mi ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Lysosome membrane ; Multi-pass membrane protein . Late endosome .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Lysosome membrane ; Multi-pass membrane protein . Late endosome .  Strongly expressed in kidney, liver, lung and pancreas.  PTM:Phosphorylated upon DNA damage, probably by ATM or
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Lysosome membrane ; Multi-pass membrane protein . Late endosome .  Strongly expressed in kidney, liver, lung and pancreas.  PTM:Phosphorylated upon DNA damage, probably by ATM or
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Lysosome membrane ; Multi-pass membrane protein . Late endosome .  Strongly expressed in kidney, liver, lung and pancreas.  PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the TMEM192 family.,

## **Products Images**



Western Blot analysis of various cells using TM192 mouse mAb