


## MBLC1 mouse mAb

Catalog No	YP-mAb-11920
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	MBLAC1
Protein Name	MBLC1
Immunogen	Synthesized peptide derived from human MBLC1 AA range: 1-51
Specificity	This antibody detects endogenous levels of MBLC1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Concentration	, mg/m
Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Cytoplasm, cytosol . Nucleus . Localizes in the nucleus during early S-phase of
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Cytoplasm, cytosol . Nucleus . Localizes in the nucleus during early S-phase of
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year  Cytoplasm, cytosol . Nucleus . Localizes in the nucleus during early S-phase of the cell cycle  cofactor:Binds 2 zinc ions per subunitsimilarity:Belongs to the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year  Cytoplasm, cytosol . Nucleus . Localizes in the nucleus during early S-phase of the cell cycle  cofactor:Binds 2 zinc ions per subunitsimilarity:Belongs to the
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Cytoplasm, cytosol . Nucleus . Localizes in the nucleus during early S-phase of the cell cycle  cofactor:Binds 2 zinc ions per subunit.,similarity:Belongs to the metallo-beta-lactamase superfamily. Glyoxalase II family.,

## **Products Images**



Western Blot analysis of various cells using MBLC1 mouse mAb