



PLOD1 mouse mAb

Catalog No	YP-mAb-12046
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	PLOD1 LLH PLOD
Protein Name	PLOD1
Immunogen	Synthesized peptide derived from human PLOD1 AA range: 551-601
Specificity	This antibody detects endogenous levels of PLOD1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Rough endoplasmic reticulum membrane; Peripheral membrane protein; Luminal side.
Tissue Specificity	
Function	catalytic activity:Procollagen L-lysine + 2-oxoglutarate + O(2) = procollagen 5-hydroxy-L-lysine + succinate + CO(2).,cofactor:Ascorbate.,cofactor:Iron.,disease:Defects in PLOD1 are the cause of Ehlers-Danlos syndrome type 6 (EDS6) [MIM:225400]. EDS is a connective tissue disorder characterized by hyperextensible skin, atrophic cutaneous scars due to tissue fragility and joint hyperlaxity. EDS6 is characterized by the presence of ocular complications, particularly retinal detachment.,disease:Defects in PLOD1 are the cause of Nevo syndrome [MIM:601451]. This is a rare, autosomal recessive disorder characterized by increased perinatal length, kyphosis, muscular hypotonia, and joint laxity. Nevo syndrome and EDS-VI have similar clinical phenotypes. Some authors consider that both syndromes are the same clinical entity.,function:Forms hydroxylysine residues in -Xaa-Lys-Gly- sequences in coll
Background	Lysyl hydroxylase is a membrane-bound homodimeric protein localized to the cisternae of the endoplasmic reticulum. The enzyme (cofactors iron and



ascorbate) catalyzes the hydroxylation of lysyl residues in collagen-like peptides. The resultant hydroxylysyl groups are attachment sites for carbohydrates in collagen and thus are critical for the stability of intermolecular crosslinks. Some patients with Ehlers-Danlos syndrome type VI have deficiencies in lysyl hydroxylase activity. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PLOD1 mouse mAb