





## PP12C mouse mAb

YP-mAb-12094
IgG
Human; Mouse
WB
PPP1R12C LENG3 MBS85
PP12C
Synthesized peptide derived from human PP12C AA range: 542-592
This antibody detects endogenous levels of PP12C at Human/Mouse
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Monoclonal, Mouse,IgG
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-1:2000
1 mg/ml
≥90%
-20°C/1 year
Cytoplasm . Cytoplasm, cytoskeleton, stress fiber .
Ubiquitously expressed. Highly expressed in heart.
caution:Although assigned as two separate genes (PPP1R12C and LENG3), it is probable that LENG3 does not exist by itself and is a part of the PPP1R12C gene.,function:Regulates myosin phosphatase activity.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Unspliced mRNA.,similarity:Contains 4 ANK repeats.,subcellular location:Along actomyosin filaments and stress fibers.,subunit:PP1 comprises a catalytic subunit, PPP1CA, PPP1CB or PPP1CC, and one or several targeting or regulatory subunits. PPP1R12C mediates binding to myosin. Interacts via its N-terminus with PPP1CB. Interacts with IL16. Interacts with the coiled-coil domain of
MPRIP.,tissue specificity:Ubiquitously expressed. Highly expressed in heart.,



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of various cells using PP12C mouse mAb